

Zeno dynamics and constraints

P Facchi^{1,5}, G Marmo², S Pascazio¹, A Scardicchio³ and
E C G Sudarshan⁴

¹ Dipartimento di Fisica, Università di Bari and Istituto Nazionale di Fisica Nucleare, Sezione di Bari, I-70126 Bari, Italy

² Dipartimento di Scienze Fisiche, Università Federico II di Napoli and Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, I-80126 Napoli, Italy

³ Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

⁴ Physics Department, University of Texas at Austin, TX 78712, USA

E-mail: paolo.facchi@ba.infn.it

Received 7 October 2003, accepted for publication 2 April 2004

Published 28 May 2004

Online at stacks.iop.org/JOptB/6/S492

DOI: 10.1088/1464-4266/6/6/006

Abstract

We investigate some examples of quantum Zeno dynamics, when a system undergoes very frequent (projective) measurements that ascertain whether it is within a given spatial region. In agreement with previously obtained results, the evolution is found to be unitary and the generator of the Zeno dynamics is the Hamiltonian with hard-wall (Dirichlet) boundary conditions. By using a new approach to this problem, this result is found to be valid in an arbitrary N -dimensional compact domain. We then propose some preliminary ideas concerning the algebra of observables in the projected region and finally look at the case of a projection onto a lower-dimensional space: in such a situation the Zeno ansatz turns out to be a procedure to impose constraints.

Keywords: quantum Zeno dynamics, constraints, decoherence, irreversibility

1. Introduction

Very frequent measurement can slow the time evolution of quantum mechanical systems. This is, in a few words, the quantum Zeno effect (QZE), by which transitions to states different from the initial one are gradually suppressed as the measurement frequency N becomes very large [1, 2] (for a review, see [3]). There are, however, two important issues that deserve attention: firstly, for a general (incomplete and nonselective [4]) measurement, represented by a complete set of projections onto multidimensional subspaces (rather than a single-dimensional one, as in the usual formulation of the QZE, by which the measurement ascertains whether the system is still in its initial, pure state), the quantum system may—and indeed does—evolve away from its initial state, although it remains in the subspace defined by the measurement (and represented by a multidimensional projection operator) [5, 6]. This leads to the formation of the ‘Zeno subspaces’ [7]. Secondly, if the measurement is not *very* frequent, the quantum evolution

yields the so-called ‘inverse’ or ‘anti-’ Zeno effect, by which transitions away from the initial state (or in general out of the relevant subspaces) are accelerated [8].

Both the Zeno and inverse Zeno phenomena have been experimentally observed during the last few years [9–12] (but see [13] for previous analyses of experimental data on nuclear hadronic cascades). The first experiment was done with an oscillating system [9], according to an interesting proposal by Cook [14], and was widely debated [15]. In a recent beautiful set of experiments, performed by Raizen’s group, first the initial quadratic and non-Markovian Zeno region was observed [10], then both the quantum Zeno and inverse Zeno effects were proved for a *bona fide* unstable system (probability leakage out of an optical potential) [11].

In this article we shall mainly analyse the first issue, investigating the features of the Zeno (sub)dynamics in the relevant subspace. This and related problems were contemplated in the seminal formulation of the QZE [2], where it was proved that the dynamics is governed by a semigroup. The details of the dynamics had interesting and

⁵ Author to whom any correspondence should be addressed.

challenging mathematical aspects, that were independently investigated by other authors [16, 17]. As a matter of fact, some mathematical issues are still unresolved nowadays. One of the most intriguing features of the original paper [2] is that some delicate operator properties were postulated on physical grounds; curiously, these postulates are always found to be valid in concrete examples, even nontrivial ones.

For a wide class of measurements, namely those represented by spatial projections, one can prove that the system evolves unitarily in a proper subspace of the total Hilbert space, the generator of the dynamics being the Hamiltonian with Dirichlet boundary conditions on the region associated with the spatial projection [5, 6]. This finding motivated further interesting studies on this topic [18–21]. In particular, Exner and Ichinose [21] analysed this result in a rigorous framework, under the nontrivial (and interesting) assumption that the original Hamiltonian be lower bounded and the Zeno Hamiltonian densely defined in the Hilbert space. The aim of this article is to further elaborate on these issues. We will first explicitly work out some examples—essentially the free case in two and three dimensions, with projections onto regular domains—and introduce a novel calculation technique, giving a constructive proof of the Zeno Hamiltonian. We then extend this result to a general spatial projection in N dimensions.

We shall prove that the Dirichlet boundary conditions are a consequence of the Zeno procedure (different proofs can be given, at different levels of generality and mathematical rigour; see [16, 5, 6, 21]), by exploring an interesting method of calculation, based on asymptotic techniques, that yields a stationary Schrödinger equation with the appropriate (Dirichlet) boundary conditions for its eigenfunctions. In section 2 we set up the general framework and introduce notation. In section 3 the projection domain is a rectangle in the plane. In section 4 it is an annulus in the plane. In section 5 we look at a spherical shell in \mathbb{R}^3 . In section 6 we generalize to regular domains in \mathbb{R}^N and in section 7 we briefly discuss the Zeno dynamics in the Heisenberg picture as well as the features of the algebra of observables in the projected domain. In section 8 we look at a different case, when the system is projected onto a domain of lower dimensionality: we shall only look at some examples and shall not attempt to generalize. One can say that in this case the Zeno ansatz yields a procedure to impose a constraint. The ideas we propose in these last two sections are somewhat embryonic and can be considered as plans for future developments. In section 9 we comment on future perspectives and applications.

2. Zeno subdynamics

Consider a free particle in N dimensions with the Hamiltonian

$$H = \frac{p^2}{2M} = -\frac{\hbar^2 \Delta}{2M}, \quad U(t) = e^{-iHt/\hbar} \quad (1)$$

acting on $\psi \in L^2(\mathbb{R}^N)$. Given a compact domain $D \subset \mathbb{R}^N$ with a nonempty interior and a regular boundary, consider the projection operator

$$P = \chi_D(x) = \int_D d^N x |x\rangle\langle x|, \quad P\psi(x) = \chi_D(x)\psi(x), \quad (2)$$

where $\chi_D(x)$ is the characteristic function of the domain D and, thought of as an operator, along with its complement $Q = 1 - P = 1 - \chi_D(x)$, decomposes the space $L^2(\mathbb{R}^N)$ into orthogonal subspaces. The Zeno subdynamics evolution operator is given by the limit

$$U_Z(t) = \lim_{N \rightarrow \infty} (G(t/N))^N, \quad (3)$$

where the (nonunitary) evolution

$$G(\tau) = PU(\tau)P \quad (4)$$

represents a single-step (projection–evolution–projection) Zeno process.

Under rather general hypotheses the limit (3) can be proved to exist [16, 2, 5, 6, 21] and yields a unitary evolution group in a proper subspace of $L^2(D)$. One gets

$$U_Z(t) = P \exp(-iH_Z t/\hbar), \quad (5)$$

where the generator of the dynamics is the Zeno Hamiltonian

$$H_Z = -\frac{\hbar^2 \Delta_D}{2M}, \quad (6)$$

defined in the domain

$$D(H_Z) = \{\psi \in L^2(D) \mid \Delta\psi \in L^2(D), \psi(\partial D) = 0\}, \quad (7)$$

∂D being the boundary of D (hard-wall or Dirichlet boundary conditions).

We will focus on this problem by looking for the eigenbasis $\{|n\rangle\}$ of $U_Z(t)$ in the subspace $PL^2(\mathbb{R}^N) \simeq L^2(D)$ such that

$$\begin{aligned} \langle n \mid U_Z(t) \mid m \rangle &= \lim_{N \rightarrow \infty} \langle n \mid G(t/N)^N \mid m \rangle = \langle n \mid e^{-iH_Z t/\hbar} \mid m \rangle \\ &= \delta_{m,n} e^{-iE_n t/\hbar}. \end{aligned} \quad (8)$$

In order to find this basis consider an arbitrary orthonormal complete set of functions in $L^2(D)$

$$\Psi_n(x) = \langle x \mid n \rangle \quad (9)$$

and take the matrix elements of the single-step operator (4)

$$G_{m,n}(t) = \langle m \mid G(t) \mid n \rangle = \text{Tr} [G(t) \mid n \rangle \langle m \mid]. \quad (10)$$

If the matrix elements of the single-step operator behave like

$$G_{m,n}(t) = \delta_{m,n} \left(1 - i \frac{E_n t}{\hbar} \right) + R_{m,n}(t), \quad (11)$$

where for $t \rightarrow 0$

$$R_{m,n}(t) = o(t), \quad (12)$$

then, under the assumption of uniform convergence of the infinite sums stemming from the insertion of $N - 1$ resolutions of the identity in (3), one obtains

$$\begin{aligned} G_{m,n}^Z(t) &\equiv \langle m \mid U_Z(t) \mid n \rangle \\ &= \lim_{N \rightarrow \infty} \sum_{n_1, \dots, n_{N-1}} G_{m,n_1}(t/N) G_{n_1,n_2}(t/N) \\ &\quad \times \dots \times G_{n_{N-1},n}(t/N) \\ &= \delta_{m,n} \exp\left(-i \frac{E_n t}{\hbar}\right). \end{aligned} \quad (13)$$

The basis $\{|n\rangle\}$ is thus the eigenbasis of H_Z belonging to the eigenvalues E_n :

$$H_Z \Psi_n(x) = E_n \Psi_n(x). \tag{14}$$

Notice that when we apply $U(t/N)$ to the relevant subspace $PL^2(\mathbb{R}^N)$, the transformed space need not be orthogonal any longer to $QL^2(\mathbb{R}^N)$, where $Q = 1 - P$, and the t/N -dependence of the scalar product of two vectors in these two subspaces is given by

$$QU(t/N)P = O(t/N). \tag{15}$$

It has been shown that equation (11) implies Dirichlet boundary conditions for the states $\Psi_n(x)$ (the ‘Zeno eigenbasis’) [6]. The proof, based on asymptotic techniques, yields the propagator in an appropriately chosen basis of eigenfunctions. In the following sections we shall introduce a novel approach: by using asymptotic analysis and the path integral representation of the matrix element (10), we will obtain a stationary Schrödinger equation and a set of boundary conditions for its eigenfunctions. This will enable us to define the induced Zeno Hamiltonian H_Z and its spectrum. The advantage of the present approach, as compared to the previous one [6], lies in the fact that one can derive a Schrödinger equation with Dirichlet boundary conditions in the projected (Zeno) subspace. Moreover, by examining some examples of multiply connected domains, we will show how the Zeno dynamics induces constraints that inherit the topological properties of the parent space.

3. Rectangle

We start off with one of the simplest examples and introduce the procedure. Consider a rectangle in the plane, $D = [0, a] \times [0, b] \subset \mathbb{R}^2$. In this case the projection (2) reads

$$P = \chi_{[0,a]}(x)\chi_{[0,b]}(y) = \int_0^a dx \int_0^b dy |xy\rangle\langle xy| \tag{16}$$

and the Hamiltonian (1) is

$$H = \frac{p_x^2 + p_y^2}{2M} = -\frac{\hbar^2}{2M}(\partial_x^2 + \partial_y^2). \tag{17}$$

The Zeno Hamiltonian, engendering the Zeno subdynamics, is formally given by (6) and (7) and represents a free particle in the box $D = [0, a] \times [0, b]$ with Dirichlet boundary conditions

$$H_Z = -\frac{\hbar^2}{2M}(\partial_x^2 + \partial_y^2), \tag{18}$$

$$\psi(0, y) = \psi(a, y) = 0, \quad \psi(x, 0) = \psi(x, b) = 0. \tag{19}$$

The eigenfunctions and eigenvalues are well known

$$\Psi_{nm}(x, y) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi}{a}x\right) \sqrt{\frac{2}{b}} \sin\left(\frac{m\pi}{b}y\right), \tag{20}$$

$$E_{nm} = \frac{\hbar^2 \pi^2}{2M} \left(\frac{n^2}{a^2} + \frac{m^2}{b^2} \right). \tag{21}$$

Let us look in detail at the derivation of the Zeno subdynamics (18)–(21) in this particular case. As explained

in section 2, the eigenbasis of the Zeno Hamiltonian H_Z in $L^2(D)$,

$$\Psi_{nm}(x, y) = \langle x, y | nm \rangle \tag{22}$$

must satisfy condition (11):

$$G_{n'm',nm}(t) = \delta_{n'n} \delta_{m'm} \left(1 - i \frac{E_{nm}t}{\hbar} \right) + o(t), \tag{23}$$

where

$$G_{n'm',nm}(t) = \langle n'm' | G(t) | nm \rangle \tag{24}$$

are the matrix elements (10) of the single-step evolution operator.

This can be proved by direct inspection: one gets

$$G_{n'm',nm}(t) = \int_0^a dx \int_0^b dy \int_0^a dx' \int_0^b dy' \left(\frac{M}{2\pi i \hbar t} \right) \times e^{i \frac{M((x'-x)^2 + (y'-y)^2)}{2\hbar t}} \Psi_{n'm'}^*(x, y) \Psi_{nm}(x', y') \tag{25}$$

and by substituting $\xi = x' - x$ and $\eta = y' - y$

$$G_{n'm',nm}(t) = \int_0^a dx \int_0^b dy \int_{-x}^{a-x} d\xi \int_{-y}^{b-y} d\eta \left(\frac{M}{2\pi i \hbar t} \right) \times e^{i \frac{M(\xi^2 + \eta^2)}{2\hbar t}} \Psi_{n'm'}^*(x, y) \Psi_{nm}(x + \xi, y + \eta). \tag{26}$$

With the natural choice $\Psi_{nm}(x, y) = \psi_n(x)\phi_m(y)$ this yields the product of two quantities

$$G_{n'm',nm}(t) = G_{n'n}(t)G_{m'm}(t) = \int_0^a dx \int_{-x}^{a-x} d\xi \left(\frac{M}{2\pi i \hbar t} \right)^{1/2} e^{i \frac{M\xi^2}{2\hbar t}} \psi_n^*(x) \psi_n(x + \xi) \times \int_0^b dy' \int_{-y}^{b-y} d\eta \left(\frac{M}{2\pi i \hbar t} \right)^{1/2} e^{i \frac{M\eta^2}{2\hbar t}} \phi_m^*(y) \phi_m(y + \eta) \tag{27}$$

and accordingly $E_{nm} = E_n + E_m$. Consider the first quantity $G_{n'n}$ and the integral over ξ . In the small- t limit there are contributions from the boundary points $\xi = -x$ and $\xi = a - x$ and from the stationary point $\xi = 0$

$$G_{n'n} = \int_0^a dx \psi_n^*(x) [\text{bound} + \text{stat}], \tag{28}$$

where

$$\text{bound} = \left(\frac{M}{2\pi i \hbar t} \right)^{1/2} \frac{\hbar t}{iM\xi} \psi_n(x + \xi) e^{i \frac{M\xi^2}{2\hbar t}} \Big|_{\xi=-x}^{\xi=a-x} + O(t^{3/2}) = \sqrt{\frac{\hbar t}{-2\pi i M}} \left(\frac{e^{iM(x-a)^2/2\hbar t}}{x-a} \psi_n(a) - \frac{e^{iMx^2/2\hbar t}}{-x} \psi_n(0) \right) + O(t^{3/2}), \tag{29}$$

while ($\lambda = M/2\hbar t$)

$$\text{stat} = \int_{-\infty}^{\infty} d\xi \sqrt{\frac{\lambda}{\pi i}} e^{i\lambda\xi^2} \times \left(\psi_n(x) + \psi_n'(x)\xi + \frac{1}{2!} \psi_n''(x)\xi^2 + O(\xi^3) \right) = \psi_n(x) + i \frac{t\hbar}{2M} \psi_n''(x) + O(t^2). \tag{30}$$

In order to obtain (23) one must require that (remember that $E_{nm} = E_n + E_m$)

$$\text{bound} = O(t^{3/2}) \quad \text{and} \quad -\frac{\hbar}{2M} \psi_n''(x) = \frac{E_n}{\hbar} \psi_n(x), \quad (31)$$

which translates into

$$-\frac{\hbar^2}{2M} \partial_x^2 \psi_n(x) = E_n \psi_n(x), \quad \text{with } \psi_n(0) = \psi_n(a) = 0, \quad (32)$$

so that for $G_{n'n}$ one obtains

$$G_{n'n}(t) = \left(1 - i \frac{E_n t}{\hbar}\right) \delta_{n'n} + O(t^{3/2}) \quad (33)$$

and analogously for $G_{m'm}$, so that

$$G_{n'm',nm}(t) = \left(1 - i \frac{E_n t}{\hbar} - i \frac{E_m t}{\hbar}\right) \delta_{n'n} \delta_{m'm} + O(t^{3/2}), \quad (34)$$

which has exactly the form (23). By equation (32) and its analogue for $\phi_m(y)$, the eigenfunctions $\Psi_{nm}(x, y) = \psi_n(x)\phi_m(y)$ of H_Z satisfy

$$-\frac{\hbar^2}{2M} (\partial_x^2 + \partial_y^2) \Psi_{n,m}(x, y) = (E_n + E_m) \Psi_{n,m}(x, y), \quad (35)$$

with Dirichlet boundary conditions. They are therefore given by (20). The Zeno Hamiltonian is therefore (18) and (19).

This derivation, although it yields the desired (and correct) result, is not rigorous. In particular, it does not tackle the delicate problem of understanding the convergence properties of the asymptotic expansion at the intersection of the (x, y) and (x', y') boundaries in equation (25) (this is apparent if one looks at the denominators of the far right-hand side of equation (29)). A similar approach will be adopted in the next sections. A more rigorous proof can be given, but will not be presented here.

4. Annulus

Consider now a circular annulus (or ring) of width $\delta r = r_2 - r_1$ on the plane, defining the domain $D = \{(x, y) | r_1^2 \leq x^2 + y^2 \leq r_2^2\}$. The projection on D reads

$$P = \chi_{[r_1, r_2]}(r) = \int_D dx dy |xy\rangle \langle xy| \\ = \int_{r_1}^{r_2} dr r \int_0^{2\pi} d\theta |r\theta\rangle \langle r\theta|. \quad (36)$$

$$r_2 - r_1 \equiv \delta r > 0. \quad (37)$$

The Zeno Hamiltonian, engendering the Zeno subdynamics, is given by (6) and represents a free particle on D with Dirichlet boundary condition

$$H_Z = -\frac{\hbar^2}{2M} (\partial_x^2 + \partial_y^2) = -\frac{\hbar^2}{2M} \left(\frac{1}{r} \partial_r (r \partial_r) + \frac{1}{r^2} \partial_\theta^2 \right), \quad (38)$$

$$\psi(r_1, \theta) = \psi(r_2, \theta) = 0. \quad (39)$$

As is well known, by writing the eigenfunctions of H_Z as $\Psi_{nl}(r, \theta) = \psi_{nl}(r)\phi_l(\theta)$, the angular functions are given by

$$\phi_l(\theta) = \frac{1}{\sqrt{2\pi}} \exp(il\theta), \quad \text{with } l = 0, \pm 1, \pm 2, \dots, \quad (40)$$

while the radial part of the eigenvalue equation reads

$$-\frac{\hbar^2}{2M} \frac{1}{r} \partial_r (r \partial_r) \psi_{nl}(r) + \frac{\hbar^2 l^2}{2M r^2} \psi_{nl}(r) = E_{nl} \psi_{nl}(r), \quad (41)$$

$$\psi_{nl}(r_1) = \psi_{nl}(r_2) = 0 \quad (42)$$

and can be solved in terms of Bessel functions.

Let us look in detail at the derivation of the Zeno subdynamics (38)–(42) in this particular case. As explained in section 2, the eigenbasis of the Zeno Hamiltonian H_Z in $L^2(D)$,

$$\Psi_{nl}(r, \theta) = \langle r, \theta | nl \rangle \quad (43)$$

must satisfy condition (11), that is

$$G_{n'l',nl}(t) = \delta_{n'n} \delta_{l'l} \left(1 - i \frac{E_{nl} t}{\hbar}\right) + o(t), \quad (44)$$

where

$$G_{n'l',nl}(t) = \langle n'l' | G(t) | nl \rangle \quad (45)$$

are the matrix elements (10) of the single-step evolution operator.

By writing $\Psi_{nl}(r, \theta) = \psi_{nl}(r)\phi_l(\theta)$, we get

$$G_{n'l',nl}(t) = \int_{r_1}^{r_2} r dr \int_{r_1}^{r_2} r' dr' \int_0^{2\pi} d\theta \int_0^{2\pi} d\theta' \\ \times \psi_{nl}(r)\phi_l(\theta)\psi_{n'l'}^*(r')\phi_{l'}^*(\theta') \left(\frac{M}{2\pi i \hbar t}\right) e^{i \frac{M d^2}{2\hbar t}}, \quad (46)$$

where d is the distance between the points (r, θ) and (r', θ')

$$d^2 = r'^2 + r^2 - 2r'r \cos(\theta' - \theta) \\ = (r' - r)^2 + 2r'r(1 - \cos(\theta' - \theta)). \quad (47)$$

Let us look first at the θ integrals. Changing again to $\eta = \theta - \theta'$ and dropping the prime one gets

$$G_{n'l',nl}(t) = \int_{r_1}^{r_2} r dr \int_{r_1}^{r_2} r' dr' \psi_{nl}(r)\psi_{n'l'}^*(r') \left(\frac{M}{2\pi i \hbar t}\right)^{1/2} \\ \times e^{i \frac{M(r-r')^2}{2\hbar t}} \int_0^{2\pi} d\theta \int_{-\theta}^{2\pi-\theta} d\eta \phi_l(\theta + \eta)\phi_{l'}^*(\theta) \\ \times \left(\frac{M}{2\pi i \hbar t}\right)^{1/2} e^{i \frac{M r'}{\hbar t} (1 - \cos \eta)}. \quad (48)$$

Consider the integral over η (at fixed r' and r). In the limit $t \rightarrow 0$ the boundary contribution reads ($z = M r' r / \hbar t$)

$$\text{bound} = \frac{1}{\sqrt{r'r}} \frac{-i}{\sqrt{2\pi i z \sin^2 \theta}} e^{iz(1 - \cos(\theta))} [\phi_l(2\pi) - \phi_l(0)] \\ + O(t^{3/2}). \quad (49)$$

In order that $O(\sqrt{t}) = O(z^{-1/2})$ vanishes and (44) is satisfied, one must require the periodicity

$$\phi_l(0) = \phi_l(2\pi). \quad (50)$$

The difference from the preceding case is given by the periodicity of the Green function.

However, we now have *two* stationary points in the η integral. One is $\eta = 0$ and the other is $\eta = \pi$ for $\theta < \pi$, or $\eta = -\pi$ for $\theta > \pi$. These represent the minimum and maximum of the distance between two points, one fixed on the circle $r' = \text{constant}$ and the other one located on the circle $r = \text{constant}$ at an angle η . One should get (at least) two points

of stationary phase each time one constrains the dynamics on a closed (hyper)surface. Both contributions must be taken into account. The only difference from the previous case is that one must also consider η^4 terms arising from the cosine in the integral

$$\text{stat}_0 = \frac{1}{\sqrt{r'r}} \int d\eta \sqrt{\frac{z}{2\pi i}} \left[\phi_l(\theta) + \frac{1}{2!} \phi_l''(\theta) \eta^2 - i \frac{z}{4!} \eta^4 \phi_l(\theta) \right] e^{iz\eta^2/2}, \quad (51)$$

$$\text{stat}_{\pm\pi} = \frac{1}{\sqrt{r'r}} e^{2iz} \int d\eta \sqrt{\frac{z}{2\pi i}} \left[\phi_l(\theta \pm \pi) + \frac{1}{2!} \phi_l''(\theta \pm \pi) \eta^2 + i \frac{z}{4!} \eta^4 \phi_l(\theta \pm \pi) \right] e^{-iz\eta^2/2}. \quad (52)$$

Notice that $\text{stat}_{\pm\pi}$ has a phase $2z = 2mr'r/\hbar t$. This phase changes the term $m(r' - r)^2/2\hbar t$ of the r', r integrals into a term $m(r' + r)^2/2\hbar t$. This factor has no more stationary points in the r', r integrals, so that its contribution can be neglected (in the $t \rightarrow 0$ limit). In turn, the contribution from $\text{stat}_{\pm\pi}$ can also be neglected. On the other hand, the stat_0 contribution is

$$\text{stat}_0 = \frac{1}{\sqrt{r'r}} \left[\phi_l(\theta) + \frac{i\hbar t}{2Mr'r} \phi_l''(\theta) + \frac{i\hbar t}{8Mr'r} \phi_l(\theta) \right] + O(t^2) \quad (53)$$

and following the same reasoning as in section 2 (rectangle on the plane) one obtains a differential equation for the eigenfunctions

$$-\phi_l''(\theta) = \alpha_l \phi_l(\theta), \quad \phi(0) = \phi(2\pi) \quad (54)$$

which yields $\alpha_l = l^2$, whence

$$\int_0^{2\pi} d\theta \phi_l^*(\theta) \text{stat}_0 = \delta_{l'l} \frac{1}{\sqrt{r'r}} \left[1 - it \frac{\hbar}{2Mr'r} \left(l^2 - \frac{1}{4} \right) \right] + O(t^2). \quad (55)$$

Therefore, the integral over r', r reads

$$\int_{r_1}^{r_2} r dr \int_{r_1}^{r_2} r' dr' \sqrt{\frac{M}{2\pi i \hbar t}} \psi_{n'l'}^*(r') \psi_{nl}(r) e^{i \frac{M(r'-r)^2}{2\hbar t}} \delta_{l'l} \times \frac{1}{\sqrt{r'r}} \left[1 - it \frac{\hbar}{2Mr'r} \left(l^2 - \frac{1}{4} \right) \right]. \quad (56)$$

By inserting $\xi = r - r'$ and dropping the prime on r' we get

$$\int_{r_1}^{r_2} \sqrt{r} dr \int_{r_1-r}^{r_2-r} \sqrt{\xi+r} d\xi \sqrt{\frac{M}{2\pi i \hbar t}} \psi_{n'l'}^*(r) \psi_{nl}(r+\xi) \times e^{i \frac{M\xi^2}{2\hbar t}} \delta_{l'l} \left[1 - it \frac{\hbar}{2Mr(r+\xi)} \left(l^2 - \frac{1}{4} \right) \right]. \quad (57)$$

By the same reasoning as before one obtains a differential equation and the Dirichlet boundary conditions for the functions $A_{nl}(r) = \sqrt{r} \psi_{nl}(r)$:

$$-\frac{\hbar^2}{2M} A_{nl}''(r) + \frac{\hbar^2}{2Mr^2} \left(l^2 - \frac{1}{4} \right) A_{nl}(r) = E_{nl} A_{nl}(r), \quad (58)$$

$$A_{nl}(r_1) = A_{nl}(r_2) = 0. \quad (59)$$

In terms of the radial functions ψ_{nl} , equation (58) becomes just equation (41), whence the Zeno Hamiltonian is given by (38) and (39).

It is interesting to notice that in this case of multiple connectedness the Zeno dynamics yields no Aharonov–Bohm topological phases. In a few words, one might say that the projected dynamics on the annulus ‘inherits’ the topological properties of the initial space \mathbb{R}^2 , and in particular the single valuedness of the wavefunction. The spatial projections do not introduce any additional ‘twist’ into the system that could induce a phase.

Two additional quick comments: first, the $r_1 \rightarrow 0$ limit yields a circle; however, it does *not* yield the Zeno dynamics on the domain $D = \{(x, y) | x^2 + y^2 < r_2^2\}$, because of the spurious condition $\psi_{nl}(0) = 0$, excluding s-wave eigenfunctions. This seemingly trivial remark clarifies that taking a limit of the projected domain does *not* necessarily yield the right Zeno dynamics. Second, the circular ring sector $\{(r, \theta) | r_1 \leq r \leq r_2, \theta_1 \leq \theta \leq \theta_2\}$ can be easily computed and yields the right dynamics and eigenfunctions (Bessel functions $I_\mu(r)$, $\mu \in \mathbb{R}$) [22].

5. Spherical shell

Let us now consider an example in \mathbb{R}^3 . We first observe that the parallelepiped can be easily dealt with by extending the techniques of section 3. We therefore look at a more interesting situation. Consider a spherical shell in \mathbb{R}^3 and a domain $D = \{(x, y, z) | r_1^2 \leq x^2 + y^2 + z^2 \leq r_2^2\}$. The projection on D reads

$$P = \chi_{[r_1, r_2]}(r) = \int_D dx dy dz |xyz| \langle xyz | \int_{r_1}^{r_2} r^2 dr \int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\phi |r\theta\phi\rangle \langle r\theta\phi|. \quad (60)$$

The Zeno Hamiltonian, engendering the Zeno subdynamics, is given by (6) and represents a free particle in the spherical shell D with Dirichlet boundary condition

$$H_Z = -\frac{\hbar^2}{2M} (\partial_x^2 + \partial_y^2 + \partial_z^2) = -\frac{\hbar^2}{2Mr^2} \left(\partial_r (r^2 \partial_r) + \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \partial_\phi^2 \right), \quad (61)$$

$$\psi(r_1, \theta, \phi) = \psi(r_2, \theta, \phi) = 0. \quad (62)$$

As is well known, by writing the eigenfunctions of H_Z as $\Psi_{nlm}(r, \theta, \phi) = R_{nl}(r) Y_{lm}(\theta) \Phi_m(\phi)$, the radial part of the eigenvalue equation reads

$$-\frac{\hbar^2}{2M} \frac{1}{r^2} \partial_r (r^2 \partial_r) R_{nl}(r) + \frac{\hbar^2}{2M} \frac{l(l+1)}{r^2} R_{nl}(r) = E_{nl} R_{nl}(r), \quad (63)$$

$$R_{nl}(r_1) = R_{nl}(r_2) = 0 \quad (64)$$

and can be solved in terms of spherical Bessel functions.

Let us see how one can obtain H_Z in this case. The first steps of the derivation are the same as before. By rewriting the distance $d(r'\theta'\phi', r\theta\phi)$ as

$$d^2 = (r' - r)^2 + 2r'r(1 - \cos(\theta' - \theta)) + 2r'r \sin \theta' \sin \theta (1 - \cos(\phi' - \phi)) \quad (65)$$

it is apparent that the integrals must be performed in the order $\phi \rightarrow \theta \rightarrow r$ and that only those stationary points that do not give an additional phase contribute to the final result.

As eigenfunctions we choose the orthogonal set

$$\Psi_{nlm}(r\theta\phi) = R_{nl}(r)Y_{lm}(\theta)\Phi_m(\phi). \quad (66)$$

The transition element is

$$\begin{aligned} G_{n'l'm',nml}(t) &= \int_{r_1}^{r_2} r'^2 dr' \int_{r_1}^{r_2} r^2 dr R_{n'l'}^*(r')R_{nl}(r) \\ &\times \sqrt{\frac{M}{2\pi\hbar t}} e^{i\frac{M(r'-r)^2}{2\hbar t}} \frac{1}{r'r} G_{l'm',lm}, \quad (67) \\ G_{l'm',lm} &= \int_0^\pi \sin\theta' d\theta' \int_0^\pi \sin\theta d\theta Y_{l'm'}^*(\theta')Y_{lm}(\theta) \\ &\times \sqrt{\frac{Mr'r}{2\pi\hbar t}} e^{i\frac{Mr'r}{\hbar}(1-\cos(\theta'-\theta))} \\ &\times \frac{1}{\sqrt{\sin\theta'\sin\theta}} \int_0^{2\pi} d\phi' \int_0^{2\pi} d\phi \sqrt{\frac{Mr'\sin\theta'r\sin\theta}{2\pi\hbar t}} \\ &\times e^{i\frac{Mr'r}{\hbar}\sin\theta'\sin\theta(1-\cos(\phi'-\phi))} \Phi_{m'}^*(\phi')\Phi_m(\phi). \quad (68) \end{aligned}$$

The ϕ' , ϕ integral is immediately computed as in the case of the annulus, section 4. Φ_m must therefore satisfy the differential equation

$$-\Phi_m'' = \alpha_m \Phi_m, \quad \Phi_m(0) = \Phi_m(2\pi), \quad (69)$$

so that $\alpha_m = m^2$. Then $G_{l'm',lm}$ becomes

$$\begin{aligned} G_{l'm',lm} &= \int_0^\pi \sqrt{\sin\theta'} d\theta' \int_0^\pi \sqrt{\sin\theta} d\theta Y_{l'm'}^*(\theta')Y_{lm}(\theta) \\ &\times \sqrt{\frac{Mr'r}{2\pi\hbar t}} e^{i\frac{Mr'r}{\hbar}(1-\cos(\theta'-\theta))} \\ &\times \left(1 - i\frac{\hbar t}{2Mr'\sin\theta'r\sin\theta}(m^2 - 1/4)\right) \delta_{m'm}. \quad (70) \end{aligned}$$

The integral over θ' , θ can be computed in a standard way (do not forget the ξ^4 term in the cosine series) and this in turn requires that the function $A_{lm} = \sqrt{\sin\theta}Y_{lm}$ must satisfy the differential equation

$$A_{lm}'' + \frac{1}{4}A_{lm} - \frac{m^2 - 1/4}{\sin^2\theta}A_{lm} = -\alpha_{lm}A_{lm}, \quad (71)$$

or, equivalently,

$$\frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial}{\partial\theta} \right) Y_{lm} - \frac{m^2}{\sin^2\theta} Y_{lm} = -\alpha_{lm} Y_{lm}, \quad (72)$$

with $Y_{lm}(0) = Y_{lm}(\pi)$. This is the standard equation for spherical harmonics. It is known that $\alpha_{lm} = l(l+1)$ irrespectively of the value of m . We obtain

$$\begin{aligned} G_{n'l'm',nml} &= \int_{r_1}^{r_2} r' dr' \int_{r_1}^{r_2} r dr R_{n'l'}^*(r')R_{nl}(r) \sqrt{\frac{M}{2\pi\hbar t}} \\ &\times e^{i\frac{M(r'-r)^2}{2\hbar t}} \left(1 - i\frac{\hbar l(l+1)}{2Mr'r} t\right) \delta_{l'l} \delta_{m'm}. \quad (73) \end{aligned}$$

Finally, the differential equation for $A_{nl} = rR_{nl}$ reads (here $E_{nl} = \frac{\hbar^2 k_{nl}^2}{2M}$, which is independent of m)

$$-A_{nl}'' + \frac{l(l+1)}{r^2} A_{nl} = k_{nl}^2 A_{nl}, \quad A_{nl}(r_1) = A_{nl}(r_2) = 0, \quad (74)$$

or, equivalently, in terms of R_{nl} , equation (63). The Zeno Hamiltonian is therefore given by (61) and (62).

6. The general case

By looking at the preceding examples one might think that the method introduced in this article is parochial and works only, for example, when the domain, besides being sufficiently regular, is also endowed with particular symmetries (regular polygons, circles, spheres and so on), that enable one to introduce coordinates with a range of integration that can be reduced to a product of intervals. In turn, this might appear as an implicit condition of separability, e.g. in the case of the three-dimensional Schrödinger equation [23]. On the contrary, as will be shown in this section, the method we propose is of general applicability.

Consider again the Hamiltonian (1) and the projection (2), $D \subset \mathbb{R}^N$ being a compact domain with nonempty interior and a regular boundary. The N -dimensional propagator (10) reads

$$\begin{aligned} G_{m,n}(t) &= \langle \mathbf{m} | G(t) | \mathbf{n} \rangle \\ &= \int_D d^N x \int_D d^N y \left(\frac{M}{2\pi\hbar t} \right)^{N/2} e^{i\frac{M(x-y)^2}{2\hbar t}} \Psi_m^*(x) \Psi_n(y) \quad (75) \end{aligned}$$

and by substituting $\xi = \mathbf{y} - \mathbf{x}$ one gets

$$\begin{aligned} G_{m,n}(t) &= \int_D d^N x \Psi_m^*(x) \int_{D-x} d^N \xi \left(\frac{M}{2\pi\hbar t} \right)^{N/2} \\ &\times e^{i\frac{M\xi^2}{2\hbar t}} \Psi_n(x+\xi) \\ &= \int_D d^N x \Psi_m^*(x) [\text{bound} + \text{stat}], \quad (76) \end{aligned}$$

where

$$D - x = \{\mathbf{y} | \mathbf{x} + \mathbf{y} \in D\}. \quad (77)$$

Let us evaluate separately the two contributions in the small- t limit. In order to compute the boundary term, we first observe that

$$e^{i\lambda\xi^2} = \frac{\xi \cdot \nabla e^{i\lambda\xi^2}}{2i\lambda\xi^2} \quad (78)$$

and then integrate by parts ($\lambda = M/2\hbar t$)

$$\begin{aligned} \text{bound} &= \int_D d^N \xi \left(\frac{\lambda}{\pi i} \right)^{N/2} \Psi_n(x+\xi) \frac{\xi \cdot \nabla e^{i\lambda\xi^2}}{2i\lambda\xi^2} \\ &= \left(\frac{\lambda}{\pi i} \right)^{N/2} \left[\int_D d^N \xi \nabla \cdot \left(\frac{\Psi_n(x+\xi)\xi e^{i\lambda\xi^2}}{2i\lambda\xi^2} \right) \right. \\ &\quad \left. - \int_D d^N \xi \nabla \cdot \left(\frac{\Psi_n(x+\xi)\xi}{\xi^2} \right) \frac{\xi \cdot \nabla e^{i\lambda\xi^2}}{(2i\lambda)^2 \xi^2} \right] \\ &= \left(\frac{\lambda}{\pi i} \right)^{N/2} \left[\oint_{\partial(D-x)} d^{N-1} S \frac{\Psi_n(x+\xi)\xi \cdot \hat{\mathbf{u}}}{\xi^2} \right. \\ &\quad \left. \times \frac{e^{i\lambda\xi^2}}{2i\lambda} (1 + O(\lambda^{-1})) \right] \\ &= \left(\frac{M}{2\pi\hbar t} \right)^{N/2} \left[\oint_{\partial D} d^{N-1} S \frac{\Psi_n(\mathbf{y})(\mathbf{y}-\mathbf{x}) \cdot \hat{\mathbf{u}}}{(\mathbf{y}-\mathbf{x})^2} \right. \\ &\quad \left. \times \frac{e^{iM(x-y)^2/2\hbar t}}{iM/\hbar t} (1 + O(t)) \right], \quad (79) \end{aligned}$$

$\hat{\mathbf{u}}$ being the unit vector perpendicular to the boundary. In the second equality, equation (78) was used again in order to obtain a higher-order volume integral with the same structure as the

initial one. The stationary contribution is obtained, as usual, by expanding the integrand function around x

$$\text{stat} = \left(\frac{M}{2\pi i\hbar t}\right)^{N/2} \int d^N \xi e^{i\lambda \xi^2} \left(\Psi_n(x) + \nabla \Psi_n(x) \cdot \xi + \frac{1}{2!} \partial_i \partial_j \Psi_n(x) \xi_i \xi_j + O(|\xi|^3) \right). \quad (80)$$

Observe that the contributions of the linear and quadratic (with $i \neq j$) terms in the integral vanish due to symmetry and one is left with

$$\text{stat} = \Psi_n(x) + i \frac{t\hbar}{2M} \Delta \Psi_n(x) + O(t^2). \quad (81)$$

In order to obtain (11) and (12) from (76) one must require that the leading contribution in the boundary term (79) vanishes and

$$-\frac{\hbar}{2M} \Delta \Psi_n(x) = \frac{E_n}{\hbar} \Psi_n(x), \quad (82)$$

namely

$$-\frac{\hbar^2}{2M} \Delta \Psi_n(x) = E_n \Psi_n(x), \quad \text{with } \Psi_n(\partial D) = 0. \quad (83)$$

Notice that the Schrödinger equation is obtained from the stationary contribution to the asymptotic expansion, while the Dirichlet boundary conditions are a consequence of the requirement that the boundary term (79) vanish at the lowest order in the expansion.

Let us briefly comment on the features of the method introduced. As already emphasized at the end of section 3, this analysis, although not entirely rigorous, yields the correct result. We derived the desired properties of the propagator by requiring at the same time the validity of the Schrödinger equation and the Dirichlet (hard-wall) boundary conditions for the eigenbasis of the (Zeno) Hamiltonian. We should emphasize, however, that the boundary and stationary terms are being dealt with separately. In fact, we did not consider the contribution of those boundary points that are *also* stationary points. Such points belong to the intersection of the boundaries of the two domains D in (75), namely $x = y \in \partial D$, and should be analysed with great care. A more rigorous treatment can be given, in which the contribution of the integral (76) is uniformly estimated: this analysis requires a different evaluation of the boundary terms and will be presented elsewhere.

The introduction of a potential [6] is not difficult to deal with if the detailed features of the convergence (10) and (11) are not worked out. Much additional care is required at a deeper mathematical level, when the self-adjointness of the Hamiltonian is called into question and must be explicitly proved. If additional rigorous results [2, 16, 17, 21] are taken into account and, by an educated guess, extended to the case of a sufficiently regular potential, one is tempted to assume that the procedure sketched above is valid in general and the Zeno dynamics governed by a self-adjoint generator (and a unitary group). The situation may clearly become more complicated when the potential is singular and/or the projected spatial region (or its boundary) lacks the required regularity.

7. Zeno dynamics in Heisenberg picture

In this section we would like to consider the Zeno dynamics in the framework of the Heisenberg picture. The following discussion must be considered preliminary: additional details and a broader picture will be given in a forthcoming paper. An interesting and natural question concerns the algebra of observables after the projection. This is not a simple problem. One can assume that to a given observable \mathcal{O} before the Zeno projection procedure there corresponds the observable POP in the projected space:

$$\mathcal{O} \Rightarrow POP. \quad (84)$$

For example, if one starts in \mathbb{R} and projects over a finite interval $P = \chi_I(x)$ (I being an interval of \mathbb{R}), the momentum and position operators become

$$p \Rightarrow PpP = \begin{cases} i\partial_x & \text{for } x \in I \\ 0 & \text{otherwise,} \end{cases} \quad (85)$$

$$x \Rightarrow PxP = \begin{cases} x & \text{for } x \in I \\ 0 & \text{otherwise.} \end{cases} \quad (86)$$

In this respect it is easy to see that the correspondence (84) is *not* an algebra homomorphism. However, if we redefine a new associative product in the algebra of operators, by setting

$$A * B \equiv APB, \quad (87)$$

with this new product the previous correspondence (84) becomes an algebra homomorphism [24]. Notice also that the new (projected) algebra acquires a unity operator P .

Notice that in general the evolution will not be an automorphism of the new product. However, it will respect the product to order $O(t/N)$ and induce, in the limit, a Zeno dynamics on the projected algebra, i.e. on the image of the projection. The evolution will be trivially an automorphism when it commutes with P and is therefore compatible with the new product without any approximation. For instance, this would be the case if we take as Hamiltonian the square of the angular momentum in the case of the annulus (section 4).

In general one has to modify the associative product in such a way that the ‘deviation’ of $U(t/N)$ from being an automorphism is of order $o(t/N)$, so that in the limit $U_Z(t)$ will be an automorphism of the new associative product adapted to the constraint. In other words, the sequence of evolution operators

$$V_N(t) = G(t/N)^N = (PU(t/N)P)^N, \quad (88)$$

yielding the Zeno limit (3), is mirrored at the level of the algebra by the following sequence of deformed associative products:

$$A *_N B \equiv AP_N B, \quad (89)$$

where P_N is a smooth positive operator with $0 \leq P_N \leq 1$ and $P_N P = P P_N = P$. For any N , P_N forms together with $Q_N = 1 - P_N$ a positive operator valued measure, yielding a resolution of the identity, i.e. $P_N + Q_N = 1$, which approximates the orthogonal resolution $P + Q = 1$, in the sense that

$$P_N \psi = P \psi + O(1/N), \quad \forall \psi \in L^2(\mathbb{R}^N). \quad (90)$$

For any N the evolution $V_N(t)$ is an automorphism of the product $*_N$ and in the limit $N \rightarrow \infty$ we get the desired result (87).

Observe that, for unbounded operators, (84) does not necessarily yield self-adjoint operators: for example, after the Zeno procedure, the momentum p would act on functions that vanish on the boundary of I and would have deficiencies $\langle 1, 1 \rangle$, see [5]. On the other hand the Zeno Hamiltonian (6) is self-adjoint. However, it would be arbitrary to require a similar property for every observable in the algebra. We shall analyse this issue in greater detail in a future article. In general, the lack of self-adjointness of the operators representing the ‘observables’ of the system in the projected subspace might be related to the incompleteness of the corresponding classical field [25, 5].

8. Projections onto lower-dimensional regions: constraints

In all the situations considered so far, the projected domain always has the same dimensionality as the original space (\mathbb{R}^n). (Remember that, after equation (1), we required the projected domain D to have a nonempty interior.) However, it is interesting to ask what would happen if one projected onto a domain D' of lower dimensionality. This is clearly a more delicate problem, as one necessarily has to face the presence of divergences. It goes without saying that these divergences must be ascribed to the lower dimensionality of the projected domain and *not* directly to the convergence features of the Zeno propagator [26]. Our problem is to understand how these divergences can be cured. One way to tackle this problem is to start from a projection onto a domain $D \subset \mathbb{R}^n$ and *then* take the limit $D \rightarrow D' \subset \mathbb{R}^{n-1}$, with a Hilbert space (Zeno subspace) $L^2(D')$.

The content of this section is preliminary. We shall only sketch the main ideas and postpone a thorough analysis to a forthcoming paper, in which the physical meaning of the divergences will be spelled out in greater details.

8.1. From the rectangle to the interval

Let us first look at the case of the rectangle, investigated in section 3, and let $b \rightarrow 0$. We first notice that in order to get a sensible result one must *first* perform the Zeno limit $N \rightarrow \infty$ and *then* let $b \rightarrow 0$. In particular one must require

$$\delta t = t/N \ll \hbar/E_m = \frac{2Mb^2}{\hbar m^2}, \quad (91)$$

which has an appealing physical meaning: the time during which the particle evolves freely between two projections must be small enough that the particle remains well within the rectangle of width b . In practice, one must first set $m < m^*$, for some m^* , in order to obtain a sensible result and finally let m^* become arbitrarily large. The order in which the two limits ($N \rightarrow \infty$ and $b \rightarrow 0$) are taken is relevant and significant from a physical perspective: one must first make sure that the wavefunction does not ‘leak’ out of the projected (Zeno) region and then let this region ‘shrink’ into a domain of lower dimensionality.

However, even if one follows the correct procedure (i.e., first $N \rightarrow \infty$ and then $b \rightarrow 0$) one still gets divergences in the phases, since

$$E_m = \frac{\hbar^2 \pi^2 m^2}{2Mb^2} \rightarrow \infty \quad \text{for } b \rightarrow 0. \quad (92)$$

Notice also that since the energy differences between different m states diverge, a superselection rule arises. Different subspaces, labelled by different values of the quantum number m , remain separated (at least for low-energy processes with energies $E \ll \hbar^2/Mb^2$). This is also physically revealing.

On the basis of the above insights, we therefore propose to perform the limit $b \rightarrow 0$ by choosing a particular eigenstate $\phi_m(y)$ and considering the reduced evolution

$$\tilde{U}_Z(t) = e^{iE_m t/\hbar} \langle m | U_Z(t) | m \rangle, \quad (93)$$

which operates only on the x degree of freedom. Physically, this corresponds to the choice of a particular value of the superselection charge. Thus, the reduced propagator reads

$$\begin{aligned} G(x', x; t) &= \langle x' | \tilde{U}_Z(t) | x \rangle = e^{iE_m t/\hbar} \langle m; x' | U_Z(t) | m; x \rangle \\ &= \sum_n e^{-iE_n t/\hbar} \psi_n(x') \psi_n^*(x), \end{aligned} \quad (94)$$

where ψ_n are the eigenfunctions of the Dirichlet problem (32), and one gets

$$\tilde{H}_Z = -\frac{\hbar^2 \partial_x^2}{2M}, \quad \text{with Dirichlet b.c.} \quad (95)$$

This is just the free particle on the interval $[0, a]$, as expected. Not only can the divergence be cured; it also yields the desired result.

The framework explained in this particular example also works in more complicated circumstances. In particular, it is important to understand in which order the two limits must be computed: first one makes sure that the Zeno mechanism works efficaciously, then takes the desired limit on the domain. We consider here two other simple situations.

8.2. From the annulus to the circle

Let us now look at the annulus, investigated in section 4. We would like to recover the evolution of a particle on a circle by considering the $\delta r \rightarrow 0$ limit, while keeping $r_1 = r_2 = R$ constant. Once again, as in section 8.1, we have to face some divergences. By taking the limit one finds the approximate eigenfunctions of equation (41)

$$\psi_{nl}(r) \simeq \psi_n(r) = \sqrt{\frac{2}{R\delta r}} \sin\left(\frac{n\pi}{\delta r}(r - R)\right) \quad (96)$$

and the energies

$$E_{nl} \simeq E_n + E_l = \frac{\hbar^2 n^2 \pi^2}{2M \delta r^2} + \frac{\hbar^2}{2MR^2} \left(l^2 - \frac{1}{4}\right). \quad (97)$$

Again one finds a diverging energy which must be regularized. However, a second (finite) term appears ($-\hbar^2/8MR^2$) [27] which is not present in the usual circle quantization. We notice that different quantization methods yield different results [28].

The reduced propagator on the remaining degree of freedom θ is just

$$G(\theta', \theta; t) = e^{i\frac{\hbar^2 n^2 \pi^2}{2M\delta r^2} t} \langle n, \theta' | U_Z(t) | n, \theta \rangle = \sum_l e^{-iE_l t/\hbar} \phi_l^*(\theta') \phi_l(\theta), \quad (98)$$

which is what one expected.

8.3. From the shell to the sphere

Finally, we reconsider the spherical shell of section 5 and take the limit $\delta r \rightarrow 0$, while keeping $r_1 = r_2 = R$ constant, as in section 8.2. This yields the energies

$$E_{nl} \simeq \frac{\hbar^2 n^2 \pi^2}{2m\delta r^2} + \frac{\hbar^2 l(l+1)}{2MR^2} \quad (99)$$

and following the same regularization procedure as before we find

$$G(\theta', \phi', \theta, \phi; t) = e^{i\frac{\hbar^2 n^2 \pi^2}{2M\delta r^2} t} \langle n; \theta', \phi' | U_Z(t) | n; \theta, \phi \rangle = \sum_{lm} e^{-i\frac{\hbar^2 l(l+1)}{2MR^2} t} Y_{lm}(\theta') \Phi_m(\phi') Y_{lm}^*(\theta) \Phi_m^*(\phi), \quad (100)$$

which is the usual propagator on a sphere of radius R , whose Hamiltonian is

$$\tilde{H}_Z = \frac{L^2}{2MR^2} = -\frac{\hbar^2}{2MR^2} \left(\frac{1}{\sin\theta} \partial_\theta (\sin\theta \partial_\theta) + \frac{1}{\sin^2\theta} \partial_\phi^2 \right). \quad (101)$$

9. Concluding remarks on potential applications

We have investigated the quantum Zeno dynamics, when a free system undergoes frequent measurements that ascertain whether it is within a sufficiently regular spatial region. The evolution in the projected (Zeno) subspace is unitary and the generator of the Zeno dynamics is the Hamiltonian with hard-wall (Dirichlet) boundary conditions. In general, this procedure leads to the formation of the ‘Zeno subspaces’ [7], on whose boundaries the wavefunction must vanish (Dirichlet): this is the ultimate reason for the absence of amplitude (and probability) leakage between ‘adjacent’ subspaces.

Quantum computation [29] is one of the most promising fields of potential application of the QZE. Interactions with the environment deteriorate the purity of quantum states and represent a very serious obstacle against the preservation of quantum superpositions and entanglement over long periods of time. It is therefore of great interest to endeavour to understand whether decoherence can be controlled and eventually halted [30]: in this context, novel techniques hinging upon the quantum Zeno effect are of interest. Besides the use of quantum error correcting codes [31], the engineering of ‘decoherence-free’ subspaces is also recently being considered and widely investigated [32]. Some mechanisms are actually being proposed, based on the so-called ‘bang–bang’ evolutions and their generalization, quantum dynamical decoupling [33]. Although ‘bang–bang’ techniques in *classical* control theory have been known to engineers since long ago [34], their introduction as a quantum control and their unification with the basic ideas underlying the quantum Zeno effect are quite

recent [35]. In particular, the decoherence-free subspaces are the dynamically generated quantum Zeno subspaces [7] within which the dynamics is far from being trivial, as has been discussed in this article. It is also worth noticing that the range of applicability of these ideas is wide, as frequent interruptions and continuous coupling [36] can yield similar physical effects. This is not entirely surprising [37], in view of Wigner’s notion of ‘spectral decomposition’ [38]. However, when one considers applications of the Zeno dynamics in the context of decoherence-free subspaces, one must remember that if the measurement is not *very* frequent, the quantum evolution yields the so-called ‘inverse’ or ‘anti-’ Zeno effect, by which transitions out of the decoherence-free subspace are accelerated [8].

In conclusion, it is interesting to notice that an issue that was considered as purely academic until a few years ago has first been experimentally demonstrated and is now being considered as a possible strategy to combat decoherence, with interesting spin-offs and very practical applications.

References

- [1] von Neumann J 1955 *Mathematical Foundation of Quantum Mechanics* (Princeton, NJ: Princeton University Press)
- Beskow A and Nilsson J 1967 *Ark. Fys.* **34** 561
- Khalfin L A 1968 *JETP Lett.* **8** 65
- [2] Misra B and Sudarshan E C G 1977 *J. Math. Phys.* **18** 756
- [3] Home D and Whitaker M A B 1997 *Ann. Phys.* **258** 237
- Facchi P and Pascazio S 2001 *Progress in Optics* vol 42, ed E Wolf (Amsterdam: Elsevier) p 147
- [4] Schwinger J 1959 *Proc. Natl Acad. Sci. USA* **45** 1552
- Schwinger J 1970 *Quantum Kinetics and Dynamics* (New York: Benjamin)
- [5] Facchi P, Gorini V, Marmo G, Pascazio S and Sudarshan E C G 2000 *Phys. Lett. A* **275** 12
- [6] Facchi P, Pascazio S, Scardicchio A and Schulman L S 2002 *Phys. Rev. A* **65** 012108
- [7] Facchi P and Pascazio S 2002 *Phys. Rev. Lett.* **89** 080401
- Facchi P and Pascazio S 2003 Quantum Zeno subspaces and dynamical superselection rules *Proc. 22nd Solvay Conf.* (Singapore: World Scientific) p 251
- [8] Lane A M 1983 *Phys. Lett. A* **99** 359
- Schieve W C, Horwitz L P and Levitan J 1989 *Phys. Lett. A* **136** 264
- Kaulakys B and Gontis V 1997 *Phys. Rev. A* **56** 1131
- Thun K and Peřina J 1998 *Phys. Lett. A* **249** 363
- Facchi P and Pascazio S 2000 *Phys. Rev. A* **62** 023804
- Elattari B and Gurvitz S A 2000 *Phys. Rev. A* **62** 032102
- Kofman A G and Kurizki G 2000 *Nature* **405** 546
- Facchi P, Nakazato H and Pascazio S 2001 *Phys. Rev. Lett.* **86** 2699
- Koshino K and Shimizu A 2003 *Phys. Rev. A* **67** 042101
- [9] Itano W M, Heinzen D J, Bollinger J J and Wineland D J 1990 *Phys. Rev. A* **41** 2295
- [10] Wilkinson S R, Bharucha C F, Fischer M C, Madison K W, Morrow P R, Niu Q, Sundaram B and Raizen M G 1997 *Nature* **387** 575
- [11] Fischer M C, Gutiérrez-Medina B and Raizen M G 2001 *Phys. Rev. Lett.* **87** 040402
- [12] Balzer C, Huesmann R, Neuhauser W and Toschek P E 2000 *Opt. Commun.* **180** 115
- [13] Valanju P, Sudarshan E C G and Chiu C B 1980 *Phys. Rev. D* **21** 1304
- [14] Cook R J 1988 *Phys. Scr. T* **21** 49
- [15] Peres A and Ron A 1990 *Phys. Rev. A* **42** 5720
- Itano W H, Heinzen D J, Bollinger J J and Wineland D J 1991 *Phys. Rev. A* **43** 5168

- Inagaki S, Namiki M and Tajiri T 1992 *Phys. Lett. A* **166** 5
 Pascazio S, Namiki M, Badurek G and Rauch H 1993 *Phys. Lett. A* **179** 155
 Blanchard Ph and Jadczyk A 1993 *Phys. Lett. A* **183** 272
 Altenmüller T P and Schenzle A 1994 *Phys. Rev. A* **49** 2016
 Berry M 1995 *Fundamental Problems in Quantum Theory* ed D M Greenberger and A Zeilinger, *Ann. New York Acad. Sci.* **755** 303
 Beige A and Hegerfeldt G 1996 *Phys. Rev. A* **53** 53
 Luis A and Peřina J 1996 *Phys. Rev. Lett.* **76** 4340
- [16] Friedman C N 1972 *Indiana Univ. Math. J.* **21** 1001
 [17] Gustafson K 1983 Irreversibility questions in chemistry, quantum-counting, and time-delay *Energy Storage and Redistribution in Molecules* ed J Hinze (New York: Plenum) and references [10,12] therein
 [18] Misra B and Antoniou A 2003 Quantum Zeno effect *Proc. 22nd Solvay Conf.* (Singapore: World Scientific) p 233
 [19] Gustafson K 2003 A Zeno story *Preprint* quant-ph/0203032
 [20] Schmidt A U 2002 *J. Phys. A: Math. Gen.* **35** 7817
 Schmidt A U 2003 *J. Phys. A: Math. Gen.* **36** 1135
 [21] Exner P and Ichinose T 2003 Product formula related to quantum Zeno dynamics *Preprint* math-ph/0302060
 [22] Sommerfeld A 1949 *Partial Differential Equation in Physics* (New York: Academic) p 168
 [23] Pauling L and Wilson E B 1935 *Introduction to Quantum Mechanics* (Singapore: McGraw-Hill) p 443
 Eisenhart L P 1934 *Phys. Rev.* **45** 428
 [24] Man'ko V I, Marmo G, Zaccaria F and Sudarshan E C G 1997 *Int. J. Mod. Phys. B* **11** 1281
 Carinena J F, Grabowski J and Marmo G 2000 *Int. J. Mod. Phys. A* **15** 4797
 [25] Nelson E 1969 *Topics in Dynamics I: Flows, Mathematica Notes* (Princeton, NJ: Princeton University Press) p 114
 Zhu C and Klauder J R 1993 *Am. J. Phys.* **61** 605
 [26] Froese R and Herbst I 2001 *Commun. Math. Phys.* **220** 489 and references therein
 [27] Goldstone J and Jaffe R L 1992 *Phys. Rev. B* **45** 14100
 [28] Scardicchio A 2002 *Phys. Lett. A* **300** 7
 [29] Galindo A and Martin-Delgado M A 2002 *Rev. Mod. Phys.* **74** 347
 Bouwmeester D, Ekert A and Zeilinger A (ed) 2000 *The Physics of Quantum Information* (Berlin: Springer)
 Nielsen M A and Chuang I L 2000 *Quantum Computation and Quantum Information* (Cambridge: Cambridge University Press)
- [30] Kofman A G and Kurizki G 2002 *Phys. Rev. Lett.* **87** 270405
 Calarco T, Datta A, Fedichev P, Pazy E and Zoller P 2003 *Phys. Rev. A* **68** 012310
 Tasaki S, Tokuse A, Facchi P and Pascazio S 2004 *Int. J. Quant. Chem.* **98** 160 (quant-ph/0210129)
 [31] Shor P W 1995 *Phys. Rev. A* **52** 2493
 Calderbank A R and Shor P W 1996 *Phys. Rev. A* **54** 1098
 Steane A 1996 *Proc. R. Soc. A* **452** 2551
 Steane A 1996 *Phys. Rev. Lett.* **77** 793
- For a review, see Preskill J 1999 *Introduction to Quantum Computation and Information* ed H K Lo, S Popescu and T P Spiller (Singapore: World Scientific)
 [32] Palma G M, Suominen K A and Ekert A K 1996 *Proc. R. Soc. A* **452** 567
 Duan L M and Guo G C 1997 *Phys. Rev. Lett.* **79** 1953
 Zanardi P and Rasetti M 1997 *Phys. Rev. Lett.* **79** 3306
 Lidar D A, Chuang I L and Whaley K B 1998 *Phys. Rev. Lett.* **81** 2594
 Knill E, Laflamme R and Viola L 2000 *Phys. Rev. Lett.* **84** 2525
 For a review, see Lidar D A and Whaley K B 2003 Decoherence-free subspaces and subsystems *Irreversible Quantum Dynamics, (Lecture Notes in Physics vol 662)* ed F Benatti and R Floreanini (Berlin: Springer) p 83 (quant-ph/0301032)
 [33] Viola L and Lloyd S 1998 *Phys. Rev. A* **58** 2733
 Viola L, Knill E and Lloyd S 1999 *Phys. Rev. Lett.* **82** 2417
 Viola L, Knill E and Lloyd S 1999 *Phys. Rev. Lett.* **83** 4888
 Viola L, Knill E and Lloyd S 2000 *Phys. Rev. Lett.* **85** 3520
 Zanardi P 1999 *Phys. Lett. A* **258** 77
 Vitali D and Tombesi P 1999 *Phys. Rev. A* **59** 4178
 Vitali D and Tombesi P 2001 *Phys. Rev. A* **65** 012305
 Uchiyama C and Aihara M 2002 *Phys. Rev. A* **66** 032313
 Byrd M S and Lidar D A 2002 *Quantum Information Processing* **1** 19
 Byrd M S and Lidar D A 2003 *Phys. Rev. A* **67** 012324
 [34] Jurdjevic V 1997 *Geometric Control Theory* (Cambridge: Cambridge University Press)
 [35] Facchi P, Lidar D A and Pascazio S 2004 *Phys. Rev. A* **69** 032314
 [36] Simonius M 1978 *Phys. Rev. Lett.* **40** 980
 Peres A 1980 *Am. J. Phys.* **48** 931
 Harris R A and Stodolsky L 1982 *Phys. Lett. B* **116** 464
 Venugopalan A and Ghosh R 1995 *Phys. Lett. A* **204** 11
 Plenio M P, Knight P L and Thompson R C 1996 *Opt. Commun.* **123** 278
 Berry M V and Klein S 1996 *J. Mod. Opt.* **43** 165
 Mihokova E, Pascazio S and Schulman L S 1997 *Phys. Rev. A* **56** 25
 Luis A and Sánchez-Soto L L 1998 *Phys. Rev. A* **57** 781
 Schulman L S 1998 *Phys. Rev. A* **57** 1509
 Thun K and Peřina J 1998 *Phys. Lett. A* **249** 363
 Panov A D 1999 *Phys. Lett. A* **260** 441
 Řeháček J, Peřina J, Facchi P, Pascazio S and Miřta L 2000 *Phys. Rev. A* **62** 013804
 Facchi P and Pascazio S 2000 *Phys. Rev. A* **62** 023804
 Militello B, Messina A and Napoli A 2001 *Phys. Lett. A* **286** 369
 Luis A 2001 *Phys. Rev. A* **64** 032104
 [37] Petrosky T, Tasaki S and Prigogine I 1990 *Phys. Lett. A* **151** 109
 Petrosky T, Tasaki S and Prigogine I 1991 *Physica A* **170** 306
 Pascazio S and Namiki M 1994 *Phys. Rev. A* **50** 4582
 [38] Wigner E P 1963 *Am. J. Phys.* **31** 6