

JULY 1981

PPPL-1788
uc-20gHAMILTONIAN FIELD DESCRIPTION OF THE
ONE-DIMENSIONAL POISSON-VLASOV EQUATIONS

BY

PHILIP J. MORRISON

PLASMA PHYSICS
LABORATORYPRINCETON UNIVERSITY
PRINCETON, NEW JERSEY

This work supported by the U.S. Department of Energy
Contract No. DE-AC02-76-CH0-3073. Reproduction, trans-
lation, publication, use and disposal, in whole or in
part, by or for the United States Government is permitted.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America.

Available from:

National Technical Information Service
U. S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22151

Price: Printed Copy \$ * ; Microfische \$3.50

<u>*PAGES</u>	<u>NTIS Selling Price</u>
1-25	\$5.00
26-50	\$6.50
51-75	\$8.00
76-100	\$9.50
101-125	\$11.00
126-150	\$12.50
151-175	\$14.00
176-200	\$15.50
201-225	\$17.00
226-250	\$18.50
251-275	\$20.00
276-300	\$21.50
301-325	\$23.00
326-350	\$24.50
351-375	\$26.00
376-400	\$27.50
401-425	\$29.00
426-450	\$30.50
451-475	\$32.00
476-500	\$33.50
500-525	\$35.00
526-550	\$36.50
551-575	\$38.00
576-600	\$39.50

For documents over 600 pages, add \$1.50 for each additional 25 page increment.

Hamiltonian Field Description of the One-Dimensional

Poisson-Vlasov Equations

by

Philip J. Morrison

Princeton University, Plasma Physics Laboratory

Princeton, New Jersey 08544

Abstract

The one-dimensional Poisson-Vlasov equations are cast into Hamiltonian form. A Poisson Bracket in terms of the phase space density, as sole dynamical variable, is presented. This Poisson bracket is not of the usual form, but possesses the commutator properties of antisymmetry, bilinearity, and nonassociativity by virtue of the Jacobi requirement. Clebsch potentials are seen to yield a conventional (canonical) formulation. This formulation is discretized by expansion in terms of an arbitrary complete set of basis functions. In particular, a wave field representation is obtained.

The Vlasov equations, in Lagrangian variables, were shown by Low¹ to possess an underlying Hamilton's principle. Here we discuss the Hamiltonian formulation of the one-dimensional version of these equations in terms of the usually encountered Eulerian description.

In connection with the development of the inverse scattering method,² Gardner³ showed that the Korteweg-deVries equation is an infinite dimensional Hamiltonian system. In so doing he obtained a Poisson bracket in terms of the noncanonical variable for this system. In a similar vein we have obtained Poisson brackets for the equations that describe a perfect fluid and ideal magnetohydrodynamics,⁴ the Maxwell-Vlasov and Poisson-Vlasov equations,⁵ and the equations which describe two-dimensional vortex fluids or guiding center plasmas.⁶

In the following we discuss the one-dimensional version of the noncanonical Poisson bracket developed in Ref. 5. It is shown that the introduction of Clebsch⁷ potentials transforms the noncanonical formulation into a canonical formulation. This formulation is discretized by expanding in terms of an arbitrary complete orthonormal set of basis functions. The exact Hamiltonian has quadratic plus quartic terms. Following this, the particular basis of Fourier decomposition is developed.

The one-dimensional Poisson-Vlasov equations are the following:

$$\frac{\partial f_{\alpha}}{\partial t}(z,t) = -v \frac{\partial f_{\alpha}}{\partial x}(z,t) + \frac{e_{\alpha}}{m_{\alpha}} \frac{\partial \phi}{\partial x}(x,t) \frac{\partial f_{\alpha}}{\partial v}(z,t) \quad (1)$$

$$\frac{\partial^2 \phi}{\partial x^2}(x,t) = - \sum_{\alpha} e_{\alpha} \int f_{\alpha}(z,t) dv \quad (2)$$

where f_{α} is the phase space probability density, ϕ is the electrostatic

potential, $z \equiv (x, v)$, α designates the species, and e_α and m_α are the signed particle charge and mass respectively. The potential ϕ is defined on R and f_α on $R \times R$. [R is the interval $(-\infty, \infty)$]. Integrating Eq. (2) twice, seeking solutions with asymptotic charge neutrality and vanishing electric field, we obtain

$$\phi(x) = - \sum_{\alpha} e_{\alpha} \int K(x|x') f_{\alpha}(z') dz' ,$$

where $K(x|x') = 1/2 |x - x'|$. (For convenience, we display only the arguments necessary to avoid confusion.) Equations (1) and (2) can be written compactly in the following form which is suggestive of the equations for two-dimensional vortex motion:⁸

$$\frac{\partial f_{\alpha}}{\partial t} = - w_{\alpha} \cdot \nabla_p f_{\alpha} , \quad (3)$$

where $\nabla_p \equiv (\partial/x, \partial/v)$ and

$$w_{\alpha} = \left(v , \frac{e_{\alpha}}{m_{\alpha}} \frac{\partial}{\partial x} \sum_{\beta} e_{\beta} \int K(x|x') f_{\beta}(z') dz' \right) .$$

Observe $\nabla_p \cdot w_{\alpha} = 0$.

In Refs. 4, 5, and 6 we discuss the properties of Poisson brackets. The brackets presented are not of the usual form, but do possess the Lie algebraic properties: bilinearity, antisymmetry, and nonassociativity as determined by the Jacobi requirement. In Ref. 5 the three-dimensional version of Eq. (3) is expressed in Heisenberg form; i.e.,

$$\frac{\partial f}{\partial t} = [f_\alpha, H] \quad , \quad (4)$$

where the bracket on the right hand side is the field Poisson bracket. With slight modification we obtain the one-dimensional Poisson bracket

$$[A, B] = \sum_{\alpha} \int \frac{f_{\alpha}(z)}{m_{\alpha}} \left\{ \frac{\delta A}{\delta f_{\alpha}}, \frac{\delta B}{\delta f_{\alpha}} \right\} dz \quad , \quad (5)$$

where the quantities A and B, on which the bracket acts, are functionals such as the integral of the Hamiltonian density:

$$H[f_{\alpha}] = \sum_{\alpha} \frac{1}{2} m_{\alpha} \int v^2 f_{\alpha}(z) dz - \frac{1}{2} \sum_{\alpha, \beta} e_{\alpha} e_{\beta} \int K(x|x') f_{\alpha}(z) f_{\beta}(z') dz dz' \quad . \quad (6)$$

The braces of the integrand of Eq. (5) signify the usual particle Poisson bracket:

$$\{f, g\} = \frac{\partial f}{\partial x} \frac{\partial g}{\partial v} - \frac{\partial f}{\partial v} \frac{\partial g}{\partial x} \quad . \quad (7)$$

The functional derivative, $\delta A / \delta f_{\alpha}$, is defined by

$$\frac{d}{d\varepsilon} A[f + \varepsilon h] \Big|_{\varepsilon=0} = \int \frac{\delta A}{\delta f} h dz \quad ,$$

where h is an arbitrary, sufficiently differentiable, function of z. Observe

$$\frac{\delta H}{\delta f_{\alpha}} = \frac{1}{2} m_{\alpha} v^2 - \sum_{\beta} e_{\alpha} e_{\beta} \int K(x|x') f_{\beta}(z') dz' \equiv H_p \quad ,$$

where H_p is the Hamiltonian (energy) of a particle in the potential

$$\phi(\mathbf{x}) = - \sum_{\beta} e_{\beta} \int K(\mathbf{x}|\mathbf{x}') f_{\beta}(z') dz' .$$

Since $\delta f(z)/\delta f(z') = \delta(z-z')$, where $\delta(z)$ is the Dirac delta function, Eq. (4) with Eq. (6) is equivalent to Eq. (3).⁹

Practical exploitation of the Hamiltonian structure presented here, either numerical or analytical, requires discretization and truncation. For noncanonical Poisson brackets, which are linear in the dynamical variables of the system, we have observed that truncation destroys the Jacobi requirement. In order to rectify this situation we seek canonical variables. To this end we represent the distribution function in terms of Clebsch potentials⁷

$$f_{\alpha}(z) = \frac{1}{m_{\alpha}} \left(\frac{\partial \bar{\Phi}^{(\alpha)}}{\partial \mathbf{x}} \frac{\partial \Gamma^{(\alpha)}}{\partial \mathbf{v}} - \frac{\partial \Gamma^{(\alpha)}}{\partial \mathbf{x}} \frac{\partial \bar{\Phi}^{(\alpha)}}{\partial \mathbf{v}} \right) . \quad (8)$$

This representation is not unique; e.g., one has the gauge condition that any function $\bar{\Gamma}^{(\alpha)}$ can be added to $\Gamma^{(\alpha)}$ provided

$$\frac{\partial \bar{\Phi}^{(\alpha)}}{\partial \mathbf{x}} \frac{\partial \bar{\Gamma}^{(\alpha)}}{\partial \mathbf{v}} - \frac{\partial \bar{\Gamma}^{(\alpha)}}{\partial \mathbf{x}} \frac{\partial \bar{\Phi}^{(\alpha)}}{\partial \mathbf{v}} = 0 .$$

Substituting Eq. (8) into the equation of motion [Eq. (3)] yields

$$\left\{ \bar{\Phi}^{(\alpha)}, \frac{\partial \Gamma^{(\alpha)}}{\partial t} + \frac{\mathbf{w}_{\alpha}}{m_{\alpha}} \cdot \nabla_{\mathbf{p}} \Gamma^{(\alpha)} \right\} + \left\{ \frac{\partial \bar{\Phi}^{(\alpha)}}{\partial t} + \frac{\mathbf{w}_{\alpha}}{m_{\alpha}} \cdot \nabla_{\mathbf{p}} \bar{\Phi}^{(\alpha)}, \Gamma^{(\alpha)} \right\} = 0 ,$$

where recall the braces are defined by Eq. (7). Clearly the above is satisfied (although not in the most general manner) if $\bar{\Phi}^{(\alpha)}$ and $\Gamma^{(\alpha)}$ satisfy

the following:¹⁰

$$\begin{aligned}\frac{\partial \Phi^{(\alpha)}}{\partial t} &= - \frac{w}{\alpha} \cdot \nabla_p \Phi^{(\alpha)} \\ \frac{\partial T^{(\alpha)}}{\partial t} &= - \frac{w}{\alpha} \cdot \nabla_p T^{(\alpha)} \quad .\end{aligned}\tag{9}$$

Upon use of the chain rule for functional derivatives, the substitution [Eq. (8)] renders the bracket [Eq. (5)] canonical:¹¹ Eqs. (9) have the form

$$\begin{aligned}\frac{\partial \Phi^{(\alpha)}}{\partial t} &= \frac{\delta H}{\delta T^{(\alpha)}} \\ \frac{\partial T^{(\alpha)}}{\partial t} &= - \frac{\delta H}{\delta \Phi^{(\alpha)}} \quad ,\end{aligned}\tag{10}$$

where H is now regarded as a functional of $\Phi^{(\alpha)}$ and $T^{(\alpha)}$.

Let us now expand $\Phi^{(\alpha)}$ and $T^{(\alpha)}$ in terms of an arbitrary orthonormal complete set of basic functions $\mu_{\underline{k}}(z)$,

$$\begin{aligned}\Phi^{(\alpha)} &= \sum_{\underline{k}} \Phi_{\underline{k}}^{(\alpha)} \mu_{\underline{k}}(z) \\ T^{(\alpha)} &= \sum_{\underline{k}} T_{\underline{k}}^{(\alpha)} \mu_{\underline{k}}(z) \quad .\end{aligned}\tag{11}$$

Here $\mu_{\underline{k}}$ is doubly indexed by \underline{k} and orthonormality, is with respect to the following inner product:

$$\langle \mu_{\underline{k}} | \mu_{\underline{l}} \rangle = \int \mu_{\underline{k}}^* \mu_{\underline{l}} dz = \delta_{\underline{k}, \underline{l}} \quad .$$

(* refers to complex conjugation and $\delta_{\underline{k},\ell}$ is the Kronecker delta function.) The reality condition is assumed; e.g., in the case where $\mu_{\underline{k}}$ is complex [e.g., Eq. (15)] in both its arguments, x and v , we require $\phi_{\underline{k}}^{(\alpha)*} = \phi_{-\underline{k}}^{(\alpha)}$ and similarly for $T_{\underline{k}}^{(\alpha)}$. In case $\mu_{\underline{k}}$ is real this condition is not applicable and the inner product is symmetric. Since an arbitrary functional $A[\phi]$ obtains its $\phi_{\underline{k}}$ dependence through ϕ , it is not difficult to verify the following:

$$\frac{\partial A}{\partial \phi_{\underline{k}}} = \int \frac{\delta A}{\delta \phi} \frac{\partial \phi}{\partial \phi_{\underline{k}}} dz = \int \frac{\delta A}{\delta \phi} \mu_{\underline{k}} dz \quad (12)$$

In this expression the A on the left hand side is to be treated as a function of the $\phi_{\underline{k}}$; in the center and on the right, A retains its role as a functional of ϕ . From Eq. (12) we obtain the following expression for the expansion of a functional derivative:

$$\frac{\delta A}{\delta \phi} = \sum_{\underline{k}} \frac{\partial A}{\partial \phi_{\underline{k}}^*} \mu_{\underline{k}} \quad .$$

With this, the discretized version of Eqs. (10) are

$$\dot{\phi}_{\underline{k}}^{(\alpha)} = \frac{\partial H}{\partial T_{\underline{k}}^{(\alpha)*}}$$

$$\dot{T}_{\underline{k}}^{(\alpha)} = - \frac{\partial H}{\partial \phi_{\underline{k}}^{(\alpha)*}} \quad .$$

These equations can be symmetrized by making the substitution

$$\psi_{\underline{k}}^{(\alpha)} = 2^{-1/2} [\phi_{\underline{k}}^{(\alpha)} + i T_{\underline{k}}^{(\alpha)}] \quad (13)$$

We obtain

$$\dot{\psi}_{\underline{k}}^{(\alpha)} = -i \frac{\partial H}{\partial \psi_{\underline{k}}^{(\alpha)*}} \quad \text{and} \quad \dot{\psi}_{\underline{k}}^{(\alpha)*} = i \frac{\partial H}{\partial \psi_{\underline{k}}^{(\alpha)}} \quad (14)$$

The distribution function (density) has an expansion which is quadratic in the $\psi_{\underline{k}}^{(\alpha)}$ and $\psi_{\underline{k}}^{(\alpha)*}$, a form which is reminiscent of quantum mechanics. From Eq. (6) we observe that the Hamiltonian has terms quadratic and quartic in the $\psi_{\underline{k}}^{(\alpha)}$ and $\psi_{\underline{k}}^{(\alpha)*}$.

Assuming periodic boundary conditions on a unit phase space box, we obtain the following wave field expression for the phase space density:

$$f_{\alpha}(z) = \sum_{\underline{k}} f_{\underline{k}}^{(\alpha)} \mu_{\underline{k}}(z) \quad ,$$

where the orthonormal functions

$$\mu_{\underline{k}} \equiv \frac{1}{2\pi} e^{ik_1 x + ik_2 y} \quad (15)$$

[Here $\underline{k} \equiv (k_1, k_2)$.] The Fourier coefficients of the density are seen by Eqs. (8), (11), and (13), to be related to the "wave function" $\psi_{\underline{k}}^{(\alpha)}$ through

$$f_{\underline{k}}^{(\alpha)} = \sum_{\underline{t}=\underline{k}+\underline{\ell}} a_{\underline{t},\underline{\ell}}^{(\alpha)} \psi_{\underline{t}}^{(\alpha)} \psi_{\underline{\ell}}^{(\alpha)*} \quad (16)$$

where

$$a_{\underline{t},\underline{\ell}}^{(\alpha)} = \frac{i}{2\pi m_{\alpha}} (t_1 \ell_2 - \ell_1 t_2) \quad .$$

Substituting Eq. (16) into Eq. (6) yields the following expression for the Hamiltonian:

$$H = \sum_{\substack{\alpha \\ l_1 = k_1 \\ l_2 \neq k_2}} S_{\underline{l}, \underline{k}}^{(2)} \psi_{\underline{l}}^{(\alpha)*} \psi_{\underline{k}}^{(\alpha)} + \sum_{\substack{\alpha, \beta \\ s_2 = l_2 \\ m_2 = p_2 \\ p_1 - m_1 = l_1 - s_1 \neq 0}} \frac{e_{\alpha}}{m_{\alpha}} \frac{e_{\beta}}{m_{\beta}} S_{\underline{l}, \underline{s}, \underline{m}, \underline{p}}^{(4)} \psi_{\underline{l}}^{(\alpha)*} \psi_{\underline{s}}^{(\alpha)} \psi_{\underline{m}}^{(\beta)*} \psi_{\underline{p}}^{(\beta)}$$

where

$$S_{\underline{l}, \underline{k}}^{(2)} = \frac{i(-1)^{l_2 - k_2}}{(l_2 - k_2)^2} (k_1 l_2 - l_1 k_2)$$

and

$$S_{\underline{l}, \underline{s}, \underline{m}, \underline{p}}^{(4)} = \frac{-(l_1 s_2 - l_2 s_1)(m_1 p_2 - m_2 p_1)}{4\pi(s_1 - l_1)(m_1 - p_1)}$$

Acknowledgements

I would like to thank J. M. Greene and A. N. Kaufman for encouragement and support. Also, I am grateful to C. Oberman and R. Kulsrud for useful conversations.

This work was supported by United States Department of Energy Contract No. DE-ACO2-76-CHO3073.

REFERENCES

1. F. E. Low, Proc. Roy. Soc. A 248, 282 (1958). For a discussion of the Lagrangian formulation of the one-dimensional Vlasov equations, see G. J. Lewak, Phys. Fluids 17, 746 (1974).
2. See the now classic papers: R. Muira, J. Math. Phys. 9, 1202 (1968); R. Muira, C. Gardner and M. Kruskal, J. Math. Phys. 9, 1204 (1968); C. Su and C. Gardner, J. Math. Phys. 10, 536 (1969); M. Kruskal, R. Muira, C. Gardner, and N. Zabusky, J. Math. Phys. 11, 952 (1970); and C. Gardner, J. Greene, M. Kruskal and R. Muira, Com. Pure Appl. Math. 27, 97 (1974). Also see Ref. 3.
3. C. S. Gardner, J. Math. Phys. 12, 1548 (1971).
4. P. J. Morrison and J. M. Greene, Phys. Rev. Lett. 45, 790 (1980).
5. P. J. Morrison, Phys. Lett. 80A, 383 (1980); see also in this connection, J. E. Marsden and A. Weinstein, to be published.
6. P. J. Morrison, Princeton Plasma Phys. Lab. Report # 1783 (1981), to be published.
7. The use of potentials to represent vector fields has a long history in hydrodynamics. For a partial historical account see, D. P. Stern, Am. J. Phys. 38, 494 (1970). See also H. Lamb, Hydrodynamics, sixth edition, (Cambridge University Press, Cambridge, 1932) sec. 167 and J. Serrin, in

Handbuch der Physik, ed. by S. Flugge (Springer-Verlag, Berlin, 1959),
Vol. 8, p. 169.

8. The Poisson bracket for the Poisson-Vlasov equation is the same as that for the vortex fluid and guiding-center plasma, although the Hamiltonians differ. See Ref. 6.
9. This requires the use of the identity, $\int f\{g,h\}dz = - \int g\{f,h\} dz$, which is readily verified by integration by parts.
10. It is well-known that, since f_α is constant on characteristics, if $f_\alpha \geq 0$ at $t=0$ for all z then $f_\alpha \geq 0$ for all z and t . If $\phi^{(\alpha)}$ and $\tau^{(\alpha)}$ are chosen at $t=0$ such that f_α , as determined by Eq. (8), is nonnegative, then the dynamical equations for $\tau^{(\alpha)}$ and $\phi^{(\alpha)}$ assure that f_α solves Eqs. (3); hence nonnegativity of f_α is maintained.
11. See Ref. 6 pp. 12 and 13.

ALL CATEGORIES

R. Askew, Auburn University, Alabama
 S. T. Wu, Univ. of Alabama
 Geophysical Institute, Univ. of Alaska
 G.L. Johnston, Sonoma State Univ, California
 H. H. Kuehl, Univ. of S. California
 Institute for Energy Studies, Stanford University
 H. D. Campbell, University of Florida
 N. L. Oleson, University of South Florida
 W. M. Stacey, Georgia Institute of Technology
 Benjamin Ma, Iowa State University
 Magne Kristiansen, Texas Tech. University
 W. L. Wiese, Nat'l Bureau of Standards, Wash., D.C.
 Australian National University, Canberra
 C.N. Watson-Munro, Univ. of Sydney, Australia
 F. Cap, Inst. for Theo. Physics, Austria
 Dr.M. Heindler, Institute for Theoretical Physics
 Technical University of Graz
 Ecole Royale Militaire, Bruxelles, Belgium
 D. Palumbo, C. European Comm. B-1049-Brussels
 P.H. Sakanaka, Instituto de Fisica, Campinas, Brazil
 M.P. Bachynski, MPB Tech., Ste. Anne de Bellevue,
 Quebec, Canada
 C. R. James, University of Alberta, Canada
 T.W. Johnston, INRS-Energie, Varennes, Quebec
 H. M. Skarsgard, Univ. of Saskatchewan, Canada
 Inst. of Physics, Academia Sinica, Peking,
 People's Republic of China
 Inst. of Plasma Physics, Hefei,
 Anhwei Province, People's Republic of China
 Library, Tsing Hua Univ., Peking, People's
 Republic of China
 Zhengwu Li, Southwestern Inst. of Phys., Leshan,
 Sichuan Province, People's Republic of China
 Librarian, Culham Laboratory, Abingdon, England (2)
 A.M. Dupas Library, C.E.N.-G, Grenoble, France
 Central Res. Inst. for Physics, Hungary
 S. R. Sharma, Univ. of Rajasthan, JAIPUR-4, India
 R. Shingal, Meerut College, India
 A.K. Sundaram, Phys. Res. Lab., India
 Biblioteca, Frascati, Italy
 Biblioteca, Milano, Italy
 G. Rostagni, Univ. Di Padova, Padova, Italy
 Preprint Library, Inst. de Fisica, Pisa, Italy
 Library, Plasma Physics Lab., Gokasho, Uji, Japan
 S. Mori, Japan Atomic Energy Res. Inst., Tokai-Mura
 Research Information Center, Nagoya Univ., Japan
 S. Shioda, Tokyo Inst. of Tech., Japan
 Inst. of Space & Aero. Sci., Univ. of Tokyo
 T. Uchida, Univ. of Tokyo, Japan
 H. Yamato, Toshiba R. & D. Center, Japan
 M. Yoshikawa, JAERI, Tokai Res. Est., Japan
 Dr. Tsuneo Nakakita, Toshiba Corporation,
 Kawasaki-Ku Kawasaki, 210 Japan
 N. Yajima, Kyushu Univ., Japan
 R. England, Univ. Nacional Auto-noma de Mexico
 B. S. Liley, Univ. of Waikato, New Zealand
 S. A. Moss, Saab Univas Norge, Norway
 J.A.C. Cabral, Univ. de Lisboa, Portugal
 O. Petrus, AL.I. CUZA Univ., Romania
 J. de Villiers, Atomic Energy Bd., South Africa
 A. Maurech, Comisaria De La Energy y Recursos
 Minerales, Spain
 Library, Royal Institute of Technology, Sweden
 Cen. de Res. En Phys.Des Plasmas, Switzerland
 Librarian, Fom-Instituut Voor Plasma-Fysica,
 The Netherlands

Bibliothek, Stuttgart, West Germany
 R.D. Buhler, Univ. of Stuttgart,
 West Germany
 Max-Planck-Inst. fur Plasmaphysik,
 W. Germany
 Nucl. Res. Estab., Julich, West Germany
 K. Schindler, Inst. Fur Theo. Physik,
 W. Germany

EXPERIMENTAL
THEORETICAL

M. H. Brennan, Flinders Univ. Australia
 H. Barnard, Univ. of British Columbia, Canada
 S. Screenivasan, Univ. of Calgary, Canada
 J. Radet, C.E.N.-B.P., Fontenay-aux-Roses,
 France
 Prof. Schatzman, Observatoire de Nice,
 France
 S. C. Sharma, Univ. of Cape Coast, Ghana
 R. N. Aiyer, Laser Section, India
 B. Buti, Physical Res. Lab., India
 L. K. Chavda, S. Gujarat Univ., India
 I.M. Las Das, Banaras Hindu Univ., India
 S. Cuperman, Tel Aviv Univ., Israel
 E. Greenspan, Nuc. Res. Center, Israel
 P. Rosenau, Israel Inst. of Tech., Israel
 Int'l. Center for Theo. Physics, Trieste, Italy
 I. Kawakami, Nihon University, Japan
 T. Nakayama, Ritsumeikan Univ., Japan
 S. Nagao, Tohoku Univ., Japan
 J.I. Sakai, Toyama Univ., Japan
 S. Tjøtta, Univ. I Bergen, Norway
 M.A. Hellberg, Univ. of Natal, South Africa
 H. Wilhelmson, Chalmers Univ. of Tech.,
 Sweden
 Astro. Inst., Sonnenborgh Obs.,
 The Netherlands
 T. J. Boyd, Univ. College of North Wales
 K. Hubner, Univ. Heidelberg, W. Germany
 H. J. Kaeppler, Univ. of Stuttgart,
 West Germany
 K. H. Spatschek, Univ. Essen, West Germany

EXPERIMENTAL
ENGINEERING

B. Grek, Univ. du Quebec, Canada
 P. Lukac, Komenskeho Univ., Czechoslovakia
 G. Horikoshi, Nat'l Lab for High Energy Physics,
 Tsukuba-Gun, Japan

EXPERIMENTAL

F. J. Paoloni, Univ. of Wollongong, Australia
 J. Kistemaker, Fom Inst. for Atomic
 & Molec. Physics, The Netherlands

THEORETICAL

F. Verheest, Inst. Vor Theo. Mech., Belgium
 J. Teichmann, Univ. of Montreal, Canada
 T. Kahan, Univ. Paris VII, France
 R. K. Chhajlani, India
 S. K. Trehan, Panjab Univ., India
 T. Namikawa, Osaka City Univ., Japan
 H. Narumi, Univ. of Hiroshima, Japan
 Korea Atomic Energy Res. Inst., Korea
 E. T. Karlson, Uppsala Univ., Sweden
 L. Stenflo, Univ. of UMEA, Sweden
 J. R. Saraf, New Univ., United Kingdom