


1. First, finish the textbook problem **10.2** — calculate to one-loop order the infinite parts of all the counterterms of the pseudoscalar Yukawa theory.

Hint: the infinite part of the four-scalar amplitude  $iV(k_1, \dots, k_4)$  does not depend on the scalar's momenta, so you may calculate it for any particular  $k_1, \dots, k_4$  you like. The off-shell momenta are OK too, so let  $k_1 = k_2 = k_3 = k_4 = 0$  — this makes for a much easier calculation of the loop diagrams. Likewise, to obtain the infinite part of the one-scalar-two-fermions amplitude, you may also let all external momenta be zeros.

2. And now consider the electric charge renormalization in the scalar QED — the theory of a EM field  $A^\mu$  interacting with a charged scalar field  $\Phi$ . At the one-loop level, there are two Feynman diagrams contributing to the 1PI two-photon amplitude, namely



- (a) Evaluate the two diagrams using dimensional regularization and verify that the net amplitude has form

$$\Sigma_{1\text{loop}}^{\mu\nu}(k) = (k^2 g^{\mu\nu} - k^\mu k^\nu) \times \Pi_{1\text{loop}}(k^2) \quad (1)$$

- (b) Calculate the  $\Pi(k^2)$  due to the above diagrams, determine the  $\delta_3$  counterterm (at the one-loop level), and write down the *net*  $\Pi(k^2)$  as a function of  $k^2$ .
- (c) Finally, consider the effective coupling  $\alpha_{\text{eff}}(k^2)$  of the scalar QED at high momenta. Show that at the one-loop level,

$$\frac{1}{\alpha_{\text{eff}}(k^2)} \approx \frac{1}{\alpha(0)} - \frac{1}{12\pi} \left( \log \frac{-k^2}{m^2} - \frac{8}{3} \right). \quad (2)$$