

Please do not waste time and paper by copying the posted homework solutions or supplementary notes. If you need to use any homework result, simply reference the appropriate question or equation and go ahead. Likewise, don't re-derive anything I derived in class.

1. The first problem is about tree-level gluon scattering, $gg \rightarrow gg$.

(a) Draw all tree diagrams for this process. Use crossing symmetry to write the net amplitude as

$$\mathcal{M}(g_1^a, g_2^b, g_3^c, g_4^d) = G^{abcd} \times \mathcal{M}_s + G^{acdb} \times \mathcal{M}_t + G^{adbc} \times \mathcal{M}_u \quad (1)$$

where G^{abcd} , *etc.*, are group factors depending on the colors of the four gluons while the \mathcal{M}_s , \mathcal{M}_t , and \mathcal{M}_u amplitudes depend on their momenta and polarizations. Thanks to the crossing symmetry,

$$\mathcal{M}_s \equiv \mathcal{M}(1, 2, 3, 4), \quad \mathcal{M}_t \equiv \mathcal{M}(1, 3, 4, 2), \quad \mathcal{M}_u \equiv \mathcal{M}(1, 4, 2, 3), \quad (2)$$

for the same analytic function \mathcal{M} applied to 3 different ordering of the four gluons. (For simplicity, treat all 4 gluons as incoming, $k_1 + k_2 + k_3 + k_4 = 0$.)

(b) Show that group factor G^{abcd} has the same index symmetry as the Riemann tensor in gravity,

$$G^{abcd} = -G^{bacd} = -G^{abdc} = +G^{cdab}, \quad (3)$$

$$G^{abcd} + G^{acdb} + G^{adbc} = 0. \quad (4)$$

Eqs. (3) should be obvious (if they are not, you have a wrong G^{abcd}), but eq. (4) takes some work. To prove it, use the identity $[[X, Y], Z] + [[Y, Z], X] + [[Z, X], Y] = 0$.


(c) Sum / average the 4-gluon |amplitude|² over all the colors and show that

$$\overline{|\mathcal{M}|^2} = \frac{C^2(G)}{2 \dim(G)} \times (3|\mathcal{M}_s|^2 + 3|\mathcal{M}_t|^2 + 3|\mathcal{M}_u|^2 - |\mathcal{M}_s + \mathcal{M}_t + \mathcal{M}_u|^2). \quad (5)$$

- (d) Prove the weak Ward identity for the 4-gluon amplitude (1): If one gluon has $e^\mu \propto k^\mu$ while the other three gluons are transverse, then $\mathcal{M} = 0$.

Hint: Show that in this case $\mathcal{M}_s = \mathcal{M}_t = \mathcal{M}_u$, then use eq. (4).

- (e) Now suppose only two gluons are transverse while the other two have unphysical polarizations (longitudinal or temporal). Or rather, let the two unphysical gluons have null polarization vectors $e^2 = 0$, specifically $e_3^\mu = k_3^\mu$ for one and $e_4^\mu k_{4\mu} = 1$ for the other. Show that in this case, the 4-gluon amplitude is exactly equal to the amplitude where the unphysical gluons are replaced with a ghost and an antighost,



Finally, let's calculate the amplitudes (2) and the partial cross-section for the four transverse gluons. For simplicity, work in the center-of-mass frame and use linear polarizations for each gluon, either \parallel to the plane of scattering or \perp to it. For the set of 4 gluons there are 16 choices of such polarizations, but the symmetries forbid some combinations and relate other combinations to each other.

- (f) Spell out which polarized $gg \rightarrow gg$ processes are forbidden and which are allowed. Write down the symmetry relations between the allowed processes. How many of them are independent?
- (g) Calculate the amplitudes (2) and the partial cross-section for the simplest choice of polarizations: all 4 gluons are \perp to the scattering plane.
- (★) Optional exercise, for extra credit:
Calculate the partial cross-sections for the other independent polarizations.
Warning: such amplitudes involve much messier algebra than the all- \perp case (g), so use Mathematica or calculate them numerically as functions of the scattering angle θ . If you try to calculate them by hand, you are liable to make more algebraic mistakes than you can fix during the time available for this exam.

2. Three exams ago — [Fall 2012 midterm](#) or [Fall 2011 midterm](#), whichever you took — you saw massive EM fields in 2 + 1 dimensions,

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{m}{4} \epsilon^{\lambda\mu\nu} A_\lambda F_{\mu\nu}. \quad (6)$$

The non-abelian version of the mass term here is the *Chern–Simons term*. In terms of matrix-valued gauge fields $\mathcal{A}_\mu = gA_\mu^a T^a$, the net Lagrangian of the *topologically massive Yang–Mills theory* in 3D is

$$\mathcal{L} = \mathcal{L}_{\text{YM}} + \mathcal{L}_{\text{CS}} = -\frac{1}{2g^2} \text{tr}(\mathcal{F}^{\mu\nu} \mathcal{F}_{\mu\nu}) + \frac{k}{8\pi} \epsilon^{\lambda\mu\nu} \text{tr} \left(\mathcal{A}_\lambda \mathcal{F}_{\mu\nu} - \frac{2i}{3} \mathcal{A}_\lambda \mathcal{A}_\mu \mathcal{A}_\nu \right). \quad (7)$$

The mass of the gluons in this theory is $M = kg^2/4\pi$; note that g^2 has dimensionality of mass in 3D, so the k coefficient — called the *Chern–Simons level* — is dimensionless. In fact, k must be integer (positive, negative, or zero) to assure the gauge invariance of the e^{iS} — and hence of the path integral of the quantum theory — despite the gauge dependence of the Chern–Simons term itself.

(a) Verify invariance of the action $\int d^3x \mathcal{L}$ under the *infinitesimal* gauge transformations.

(★) Optional exercise, for extra credit:

Show that under a finite gauge transformation $U(x)$ the action changes by

$$\Delta S = \frac{-k}{12\pi} \int d^3x \epsilon^{\lambda\mu\nu} \text{tr} \left(U^{-1} \partial_\lambda U \cdot U^{-1} \partial_\mu U \cdot U^{-1} \partial_\nu U \right). \quad (8)$$

FYI — but don't try to prove this during the exam — the integral here depends only on the topological properties of the $U(x)$ and its values are always integer $\times 24\pi^2$. Consequently, $\Delta S = 2\pi k \times \text{an integer}$, hence for integer Chern–Simons levels k the e^{iS} is gauge invariant.

The rest of this exercise is about the quantum gauge theories in 3D. Specifically, we shall see how in 3D QCD, the loops of massive quarks *induce* an effective Chern–Simons term for the gluons.


For simplicity, consider the 3D $SU(N)$ gauge theory with a single fundamental multiplet \mathbf{N} of quarks (*i.e.*, N colors, one flavor) and no tree-level CS term, thus

$$\mathcal{L} = \frac{-1}{2g^2} \text{tr}(\mathcal{F}^{\mu\nu} \mathcal{F}_{\mu\nu}) + \bar{\Psi}(i\not{D} - m)\Psi. \quad (9)$$

Note that in odd spacetime dimensions, massive Dirac fermions have no Parity symmetry even at the tree level. At higher orders of perturbation theory, the quark loop diagrams

yield parity-violating amplitudes thanks to $\underline{\text{tr}(\gamma^\lambda \gamma^\mu \gamma^\nu) = 2i\epsilon^{\lambda\mu\nu}}$ in 3D (and similar formulæ in higher odd dimensions), so the gluon sector of the theory also becomes parity-violating.

(b) Evaluate the one loop diagram



and show that for *small* gluon momentum $|p| \ll m$ it yields

$$\Sigma_{\psi \text{ loop}}^{\mu a, \nu b}(p) = \frac{g^2 \delta^{ab}}{8\pi} \left(-ip_\lambda \epsilon^{\lambda\mu\nu} + \frac{p^\mu p^\nu - g^{\mu\nu} p^2}{3m} + O\left(\frac{p^3}{m^2}\right) \right). \quad (10)$$

(c) Similarly, show that for three external gluons with small momenta (compared to the fermion's mass m), the one-loop amplitude is

$$iV_{\lambda\mu\nu}^{abc} = \text{[Diagram 1]} + \text{[Diagram 2]} = \frac{ig^3}{8\pi} f^{abc} \epsilon_{\lambda\mu\nu} + O\left(\frac{p}{m}\right). \quad (11)$$

(d) Show that for quark loops with four or more external gluons with small momenta, all the one-quark-loop amplitudes are suppressed by negative powers of the quark mass m .

Now consider the Functional Integral for the $d = 3$ QCD. Let us integrate $\iint D[\Psi(x)] \iint D[\bar{\Psi}(x)]$ over the quark fields for fixed gauge fields $A_\mu^a(x)$. The result of this integration is an effective quantum theory of the gauge fields with action

$$S[A_\mu^a] = S_{YM}[A_\mu^a] - i \log \text{Det}(i\mathcal{D} - m). \quad (12)$$

(e) Use the results of questions (b–d) to show that in the large quark mass m limit,

$$-i \log \text{Det}(i\mathcal{D} - m) = \frac{1}{2} \int d^3x \left\{ \frac{1}{8\pi} \epsilon^{\lambda\mu\nu} \text{tr}(\mathcal{A}_\lambda \mathcal{F}_{\mu\nu} - \frac{2i}{3} \mathcal{A}_\lambda \mathcal{A}_\mu \mathcal{A}_\nu) + O\left(\frac{1}{m}\right) \right\} \quad (13)$$

and consequently, the effective low-energy quantum theory is precisely the topologically massive Yang–Mills theory (7) with Chern–Simons level $k = \frac{1}{2}$.

Finally, consider 3D QCD with several flavors of massive quarks, some with $m_f > 0$ and some with $m_f < 0$ (in 3D, this makes a difference). Let's also have a tree-level Chern–Simons level k_0 .

(f) Show that when we integrate out all the quarks, we end up with the net Chern–Simons level

$$k = k_0 + \frac{\#(m_f > 0) - \#(m_f < 0)}{2}. \quad (14)$$

Note: *consistency of the quantum theory requires the net CS level k to be integer*. Consequently, in theories with even n_f the tree-level k_0 should be integer, but the theories with odd n_f should have half-integer $k_0 \in \mathbf{Z} + \frac{1}{2}$.