

1. Back in 1996, Aharoni, Sonnenschein, Theisen, and Yankielowicz [arXiv:hep-th/9611222](https://arxiv.org/abs/hep-th/9611222) found a curious $SU(2) \times SU(2)$ SUSY gauge theory ‘living’ on a D3-brane probe located near an intersection of two orientifold planes. In this problem, we study the renormalization group flow and the IR fixed points of this gauge theory from a purely 4D QFT point of view.

Ignoring the ‘stringy’ degrees of freedom, we have $d = 4$, $\mathcal{N} = 1$ SUSY gauge theory with $G = SU(2) \times SU(2)$ and the following multiplets of chiral superfields:

$$A_1, A_2 \in (\mathbf{2}, \mathbf{2}), \quad B_1, \dots, B_8 \in (\mathbf{2}, \mathbf{1}), \quad C_1, \dots, C_8 \in (\mathbf{1}, \mathbf{2}). \quad (1)$$

Note: in my notations I keep explicit flavor indices but suppress the gauge indices, so the net number of chiral superfields in this theory is

$$(2 \times 2 \times 2)_A + (8 \times 2 \times 1)_B + (8 \times 1 \times 2)_C = 8 + 16 + 16 = 40.$$

The theory has a superpotential

$$W = \lambda \sum_{i=1}^4 B_i A_1 C_i + \lambda \sum_{i=5}^8 B_i A_2 C_i. \quad (2)$$

Again, the gauge indices are suppressed in this formula, but for each term here there is only one gauge-invariant way of contracting the gauge indices.

- (a) List global symmetries of this model and show that it has only 3 independent anomalous dimensions: γ_A (same for A_1 and A_2), γ_B (same for all B_i), and γ_C (same for all C_i). Allow for different gauge couplings $g_1 \neq g_2$ of the two $SU(2)$ factors.
- (b) Calculate the exact β_λ , β_{g_1} , and β_{g_2} in terms of the anomalous dimensions and show that all three beta-functions vanish when $\gamma_B = \gamma_C = -\frac{1}{2}\gamma_A$. Argue that this leads to a line of fixed points in the (λ, g_1, g_2) coupling space.

- (c) Calculate the anomalous dimensions γ_A , γ_B , and γ_C to one-loop order and show that the fixed line lies at

$$g_1^2 = g_2^2 = \frac{16}{12} \lambda^2 + O(\lambda^4). \quad (3)$$

- (d) Show that this fixed line is IR-attractive. That is, if we start with some other couplings in the UV and let the RG run to lower energies, then in the IR limit the couplings will end somewhere on the fixed line (3).
- (e) Any IR-attractive fixed point gives rise to an SCFT (super-conformal field theory), and a line (surface, *etc.*) of such fixed points makes a whole family of non-trivial SCFTs. Argue that for the model in question, this family of SCFTs includes both weakly-coupled and strongly-coupled theories.
2. In lieu of the second problem, finish reading the [1982 Witten's paper *Constraints On Supersymmetry Breaking*](#) that I have assigned last week.

Skim over section 9 of the paper — the group theory there is hard to follow for non-experts. More importantly, the main result of section 9 is not quite right, as Witten himself had clarified in his 2000 paper [arXiv:hep-th/0006010](#). Specifically, a pure super-Yang–Mills theory with gauge group G has Witten's index $I = C(G)$ — the Casimir of the adjoint multiplet — rather than $I = \text{rank}(G) + 1$. For the $SU(N)$ and $Sp(N)$ groups both formulae give the same answer, but for the $SO(N)$ and the exceptional groups there is a difference.