

1. First, consider a scalar analogue of QCD, or more generally a non-abelian gauge theory with some gauge group G and complex scalar fields $\Phi^i(x)$ in some multiplet (r) of G .

(a) Write down the physical Lagrangian of this theory, the complete bare Lagrangian of the quantum theory in the Feynman gauge, and the Feynman rules.

Now consider the annihilation process $\Phi + \Phi^* \rightarrow 2$ gauge bosons. At the tree level, there are four Feynman diagrams contributing to this process.

(b) Draw the diagrams and write down the tree-level annihilation amplitude.

As discussed in class, amplitudes involving the non-abelian gauge fields satisfy a weak form of the Ward identity: *On-shell Amplitudes involving a longitudinally polarized gauge bosons vanish, provided all the other gauge bosons are transversely polarized.* In other words,

$$\mathcal{M} \equiv e_1^{\mu_1} e_2^{\mu_2} \cdots e_n^{\mu_n} \mathcal{M}_{\mu_1 \mu_2 \cdots \mu_n}(\text{momenta}) = 0$$

when $e_1^\mu \propto k_1^\mu$ but $e_2^\nu k_{2\nu} = \cdots = e_n^\nu k_{n\nu} = 0$.

(c) Verify this identity for the scalar annihilation amplitude.

2. Next, a bit of group theory. Consider a generic simple non-abelian compact Lie group G and its generators T^a . For a suitable normalization of the generators,

$$\text{tr}_{(r)}(T^a T^b) \equiv \text{tr} \left(T_{(r)}^a T_{(r)}^b \right) = R(r) \delta^{ab} \quad (1)$$

where the trace is taken over any complete multiplet (r) — irreducible or reducible, it does not matter — and $T_{(r)}^a$ is the matrix representing the generator T^a in that multiplet. The coefficient $R(r)$ in eq. (1) depends on the multiplet (r) but it's the same for all generators T^a and T^b . The $R(r)$ is called the *index* of the multiplet (r).

The (quadratic) Casimir operator $C_2 = \sum_a T^a T^a$ commutes with all the generators, $\forall b, [C_2, T^b] = 0$. Consequently, when we restrict this operator to any *irreducible* multiplet (r) of the group G , it becomes a unit matrix times some number $C(r)$. In other words,

$$\text{for an irreducible } (r), \quad \sum_a T_{(r)}^a T_{(r)}^a = C(r) \times \mathbf{1}_{(r)}. \quad (2)$$

For example, for the isospin group $SU(2)$, the Casimir operator is $C_2 = \vec{I}^2$, the irreducible multiplets have definite isospin $I = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$, and $C(I) = I(I + 1)$.

(a) Show that for any irreducible multiplet (r),

$$\frac{R(r)}{C(r)} = \frac{\dim(r)}{\dim(G)}. \quad (3)$$

In particular, for the $SU(2)$ group, this formula gives $R(I) = \frac{1}{3}I(I + 1)(2I + 1)$.

(b) Suppose the first three generators T^1, T^2 , and T^3 of G generate an $SU(2)$ subgroup, thus

$$[T^1, T^2] = iT^3, \quad [T^2, T^3] = iT^1, \quad [T^3, T^1] = iT^2. \quad (4)$$

Show that if a multiplet (r) of G decomposes into several $SU(2)$ multiplets of isospins I_1, I_2, \dots, I_n , then

$$R(r) = \sum_{i=1}^n \frac{1}{3} I_i (I_i + 1) (2I_i + 1). \quad (5)$$

(c) Now consider the $SU(N)$ group with an obvious $SU(2)$ subgroup of matrices acting only on the first two components of a complex N -vector. This complex N -vector is called the fundamental multiplet (of the $SU(N)$) and denoted (N) or \mathbf{N} . As far as the $SU(2)$ subgroup is concerned, (N) comprises one doublet and $N - 2$ singlets, hence

$$R(N) = \frac{1}{2} \quad \text{and} \quad C(N) = \frac{N^2 - 1}{2N}. \quad (6)$$

Show that the adjoint multiplet of the $SU(N)$ decomposes into one $SU(2)$ triplet, $2(N - 2)$ doublets, and $(N - 2)^2$ singlets, therefore

$$R(\text{adj}) = C(\text{adj}) \equiv C(G) = N. \quad (7)$$

Hint: $(N) \times (\overline{N}) = (\text{adj}) + (1)$.

(d) The symmetric and the anti-symmetric 2-index tensors form irreducible multiplets of the $SU(N)$ group. Find out the decomposition of these multiplets under the $SU(2) \subset SU(N)$ and calculate their respective indices R and Casimirs C .

3. Now let's apply this group theory to physics. Consider quark-antiquark pair production in QCD, specifically $u\bar{u} \rightarrow d\bar{d}$. There is only one tree diagram contributing to this process,



Evaluate this diagram, then sum/average the $|\mathcal{M}|^2$ over both spins and *colors* of the final/initial particles to calculate the total cross section. For simplicity, you may neglect the quark masses.

Note that the diagram (8) looks exactly like the QED pair production process $e^-e^+ \rightarrow \text{virtual } \gamma \rightarrow \mu^-\mu^+$, so you can re-use the QED formula for summing/averaging over the spins, *cf.* [my notes on Dirac traceology from the Fall semester](#), page 11. But in QCD, you should also sum/average over the colors of all the quarks, and that's the whole point of this exercise.

4. Finally, let's continue problem 1 but focus on the group theory and cross-sections rather than the Ward identity.

(a) Go back to the gauge theory from problem 1 and the tree-level annihilation amplitude of a scalar 'quark' Φ^i and an 'antiquark' Φ_j^* into a pair of gauge bosons with adjoint colors a and b . Take the annihilation amplitude from part (b) of problem 1, focus on its color dependence, and rewrite it in the form

$$\mathcal{M}(j + i \rightarrow a + b) = F \times \{T^a, T^b\}_j^i + iG \times [T^a, T^b]_j^i \quad (9)$$

where F and G are some functions of all the momenta and of the vectors' polarizations. Give explicit formulae for F and G .

- (b) Next, let us sum the $|\mathcal{M}|^2$ over the gauge boson's colors and average over the scalars' colors. Show that

$$\frac{1}{\dim^2(r)} \sum_{ij} \sum_{ab} |\mathcal{M}|^2 = \frac{C(r)}{\dim(r)} \times \left((4C(r) - C(\text{adj})) \times |F|^2 + C(\text{adj}) \times |G|^2 \right). \quad (10)$$

In particular, for scalars in the fundamental representation of the $SU(N)$ gauge group,

$$\frac{1}{N^2} \sum_{ij} \sum_{ab} |\mathcal{M}|^2 = \frac{N^2 - 1}{2N^2} \left(\frac{N^2 - 2}{N} \times |F|^2 + N \times |G|^2 \right). \quad (11)$$

- (c) Evaluate F and G in the center of mass frame, where the vector particles' polarizations $e_{1,2}^\mu = (0, \mathbf{e}_{1,2})$ are purely spatial and transverse to the vectors' momenta $\pm \mathbf{k}$. For simplicity, use planar rather than circular polarizations.
- (d) Assemble your results and calculate the (polarized, partial) cross-section for the annihilation process.