

1. Consider the plane-wave solutions of the Dirac equation, $\Psi_\alpha(x) = u_\alpha \times e^{-ipx}$ and $\Psi_\alpha(x) = v_\alpha \times e^{+ipx}$ for some x -independent Dirac spinors $u_\alpha(p, s)$ and $v_\alpha(p, s)$.

- (a) Check that these waves indeed solve the Dirac equation provided $p^2 = m^2$ while

$$(\not{p} - m)u(p, s) = 0, \quad (\not{p} + m)v(p, s) = 0. \quad (1)$$

By convention, we always take $E = p^0 = +\sqrt{\mathbf{p}^2 + m^2}$ — that's why we have both $e^{-ipx}u_\alpha$ and $e^{+ipx}v_\alpha$ types of wave — while the spinor coefficients $u(p, s)$ and $v(p, s)$ are normalized to

$$u^\dagger(p, s)u(p, s') = v^\dagger(p, s)v(p, s') = 2E\delta_{s,s'}. \quad (2)$$

In this problem we shall write down explicit formulae for these spinors in the Weyl basis for the γ^μ matrices.

- (b) Show that for $\mathbf{p} = 0$,

$$u(\mathbf{p} = \mathbf{0}, s) = \begin{pmatrix} \sqrt{m} \xi_s \\ \sqrt{m} \xi_s \end{pmatrix} \quad (3)$$

where ξ_s is a two-component $SO(3)$ spinor encoding the electron's spin state. The ξ_s are normalized to $\xi_s^\dagger \xi_{s'} = \delta_{s,s'}$.

- (c) For other momenta, $u(p, s) = M_D(\text{boost}) \times u(\mathbf{p} = 0, s)$ for the boost that turns $(m, \vec{0})$ into p^μ . Use eqs. (7.19) from the [previous homework set #7](#) to show that

$$u(p, s) = \begin{pmatrix} \sqrt{E - \mathbf{p} \cdot \boldsymbol{\sigma}} \xi_s \\ \sqrt{E + \mathbf{p} \cdot \boldsymbol{\sigma}} \xi_s \end{pmatrix} = \begin{pmatrix} \sqrt{p_\mu \bar{\sigma}^\mu} \xi_s \\ \sqrt{p_\mu \sigma^\mu} \xi_s \end{pmatrix}. \quad (4)$$

- (d) Use similar arguments to show that

$$v(p, s) = \begin{pmatrix} +\sqrt{E - \mathbf{p} \cdot \boldsymbol{\sigma}} \eta_s \\ -\sqrt{E + \mathbf{p} \cdot \boldsymbol{\sigma}} \eta_s \end{pmatrix} = \begin{pmatrix} +\sqrt{p_\mu \bar{\sigma}^\mu} \eta_s \\ -\sqrt{p_\mu \sigma^\mu} \eta_s \end{pmatrix} \quad (5)$$

where η_s are two-component $SO(3)$ spinors normalized to $\eta_s^\dagger \eta_{s'} = \delta_{s,s'}$.

Physically, the η_s should have opposite spins from ξ_s — the holes in the Dirac sea have opposite spins (as well as p^μ) from the missing negative-energy particles. Mathematically, this requires $\eta_s^\dagger \mathbf{S} \eta_s = -\xi_s^\dagger \mathbf{S} \xi_s$; we may solve this condition by letting $\eta_s = \sigma_2 \xi_s^* = \pm i \xi_{-s}^*$.

- (e) Check that $\eta_s = \sigma_2 \xi_s^* = \pm i \xi_{-s}^*$ indeed provides for the $\eta_s^\dagger \mathbf{S} \eta_s = -\xi_s^\dagger \mathbf{S} \xi_s$, then show that this leads to

$$v(p, s) = \gamma^2 u^*(p, s) \quad \text{and} \quad u(p, s) = \gamma^2 v^*(p, s). \quad (6)$$

- (f) Show that for the ultra-relativistic electrons or positrons of definite helicity $\lambda = \pm \frac{1}{2}$, the Dirac plane waves become *chiral* — *i.e.*, dominated by one of the two irreducible Weyl spinor components $\psi_L(x)$ or $\psi_R(x)$ of the Dirac spinor $\Psi(x)$, while the other component becomes negligible. Specifically,

$$\begin{aligned} u(p, -\tfrac{1}{2}) &\approx \sqrt{2E} \begin{pmatrix} \xi_L \\ 0 \end{pmatrix}, & u(p, +\tfrac{1}{2}) &\approx \sqrt{2E} \begin{pmatrix} 0 \\ \xi_R \end{pmatrix}, \\ v(p, -\tfrac{1}{2}) &\approx -\sqrt{2E} \begin{pmatrix} 0 \\ \eta_L \end{pmatrix}, & v(p, +\tfrac{1}{2}) &\approx \sqrt{2E} \begin{pmatrix} \eta_R \\ 0 \end{pmatrix}. \end{aligned} \quad (7)$$

Note that for the electron waves the helicity agrees with the chirality — they are both left or both right, — but for the positron waves the chirality is opposite from the helicity.

In the [previous homework#7](#) (problem 6.g), we saw that for $m = 0$ the LH and the RH Weyl spinor fields decouple from each other. Now this exercise show us which particle modes comprise each Weyl spinor: *The $\psi_L(x)$ and its hermitian conjugate $\psi_L^\dagger(x)$ contain the left-handed fermions and the right-handed antifermions, while the $\psi_R(x)$ and the $\psi_R^\dagger(x)$ contain the right-handed fermions and the left-handed antifermions.*

2. Next, let's establish some basis-independent properties of the Dirac spinors $u(p, s)$ and $v(p, s)$ — although you may use the Weyl basis to verify them. We shall use these properties in class when we get to Quantum Electrodynamics (QED).

- (a) Show that

$$\bar{u}(p, s) u(p, s') = +2m \delta_{s, s'}, \quad \bar{v}(p, s) v(p, s') = -2m \delta_{s, s'}; \quad (8)$$

note the $\pm 2m$ normalization factors here, unlike the $+2E$ factors in eq. (2) for the $u^\dagger u$ and the $v^\dagger v$.

- (b) There are only two independent $SO(3)$ spinors, hence $\sum_s \xi_s \xi_s^\dagger = \sum_s \eta_s \eta_s^\dagger = \mathbf{1}_{2 \times 2}$. Use this fact to show that

$$\sum_{s=1,2} u_\alpha(p, s) \bar{u}_\beta(p, s) = (\not{p} + m)_{\alpha\beta} \quad \text{and} \quad \sum_{s=1,2} v_\alpha(p, s) \bar{v}_\beta(p, s) = (\not{p} - m)_{\alpha\beta}. \quad (9)$$

3. Now consider the charge conjugation symmetry \mathbf{C} which exchanges particles with antiparticles, for example the electrons e^- with the positrons e^+ ,

$$\widehat{\mathbf{C}} |e^-(\mathbf{p}, s)\rangle = |e^+(\mathbf{p}, s)\rangle, \quad \widehat{\mathbf{C}} |e^+(\mathbf{p}, s)\rangle = |e^-(\mathbf{p}, s)\rangle. \quad (10)$$

Note that the operator $\widehat{\mathbf{C}}$ is unitary and squares to one (repeating the exchange brings us back to the original particles), hence $\widehat{\mathbf{C}}^\dagger = \widehat{\mathbf{C}}^{-1} = \widehat{\mathbf{C}}$.

- (a) In the fermionic Fock space, the $\widehat{\mathbf{C}}$ operator act on multi-particle states by turning each particle into an antiparticle and vice versa according to eqs. (10). Show that this action implies

$$\widehat{\mathbf{C}} \hat{a}_{\mathbf{p},s}^\dagger \widehat{\mathbf{C}} = \hat{b}_{\mathbf{p},s}^\dagger, \quad \widehat{\mathbf{C}} \hat{b}_{\mathbf{p},s}^\dagger \widehat{\mathbf{C}} = \hat{a}_{\mathbf{p},s}^\dagger, \quad \widehat{\mathbf{C}} \hat{a}_{\mathbf{p},s} \widehat{\mathbf{C}} = \hat{b}_{\mathbf{p},s}, \quad \widehat{\mathbf{C}} \hat{b}_{\mathbf{p},s} \widehat{\mathbf{C}} = \hat{a}_{\mathbf{p},s}. \quad (11)$$

- (b) The quantum Dirac fields $\widehat{\Psi}(x)$ and $\widehat{\bar{\Psi}}(x)$ are linear combinations of creation and annihilation operators. Use eqs. (11) and the plane-wave relations (6) to show that

$$\widehat{\mathbf{C}} \widehat{\Psi}(x) \widehat{\mathbf{C}} = \gamma^2 \widehat{\bar{\Psi}}^*(x) \quad \text{and} \quad \widehat{\mathbf{C}} \widehat{\bar{\Psi}}(x) \widehat{\mathbf{C}} = \widehat{\Psi}^*(x) \gamma^2 \quad (12)$$

where $*$ stands for the hermitian conjugation of the component fields but without transposing a column vector (of 4 Dirac components) into a row vector or vice versa, thus

$$\widehat{\Psi} = \begin{pmatrix} \hat{\psi}_1 \\ \hat{\psi}_2 \\ \hat{\psi}_3 \\ \hat{\psi}_4 \end{pmatrix}, \quad \widehat{\bar{\Psi}} = \begin{pmatrix} \hat{\psi}_1^\dagger \\ \hat{\psi}_2^\dagger \\ \hat{\psi}_3^\dagger \\ \hat{\psi}_4^\dagger \end{pmatrix}, \quad \begin{aligned} \widehat{\Psi} &= (\hat{\psi}_1^\dagger, \hat{\psi}_2^\dagger, \hat{\psi}_3^\dagger, \hat{\psi}_4^\dagger) \times \gamma^0, \\ \widehat{\bar{\Psi}}^* &= (\hat{\psi}_1, \hat{\psi}_2, \hat{\psi}_3, \hat{\psi}_4) \times \gamma^0. \end{aligned} \quad (13)$$

- (c) Show that the Dirac equation transforms covariantly under the charge conjugation (12). Hint: prove and use $\gamma^\mu \gamma^2 = -\gamma^2 (\gamma^\mu)^*$ for all γ^μ in the Weyl basis.

- (d) Show that that the *classical* Dirac Lagrangian is invariant under the charge conjugation (up to a total spacetime derivative). Note that in the classical limit the Dirac fields *anticommute* with each other, $\Psi_\alpha^* \Psi_\beta = -\Psi_\beta \Psi_\alpha^*$. Also, similar to the hermitian conjugation of quantum fields, the complex conjugation of fermionic fields reverses their order: $(F_1 F_2)^* = F_2^* F_1^* = -F_1^* F_2^*$.

4. Another important discrete symmetry is the *parity* \mathbf{P} , the im-proper Lorentz symmetry that reflects the space but not the time, $(\mathbf{x}, t) \rightarrow (-\mathbf{x}, +t)$. This symmetry acts on the Dirac spinor fields according to

$$\widehat{\Psi}'(-\mathbf{x}, +t) = \pm \gamma^0 \widehat{\Psi}(\mathbf{x}, +t) \quad (14)$$

where the overall \pm sign is the *intrinsic parity* of the fermion species described by the $\widehat{\Psi}$ field.

- (a) Verify that the Dirac equation transforms covariantly under (14) and that the Dirac Lagrangian is invariant (apart from $\mathcal{L}(\mathbf{x}, t) \rightarrow \mathcal{L}(-\mathbf{x}, t)$).

In the Fock space, eq. (14) becomes

$$\widehat{\mathbf{P}} \widehat{\Psi}(\mathbf{x}, t) \widehat{\mathbf{P}} = \pm \gamma^0 \widehat{\Psi}(-\mathbf{x}, t) \quad (15)$$

for some unitary operator $\widehat{\mathbf{P}}$ that squares to one. Let's find how this operator acts on the particles and their states.

- (b) First, check the plane-wave solutions $u(\mathbf{p}, s)$ and $v(\mathbf{p}, s)$ from problem (1) and show that $u(-\mathbf{p}, s) = +\gamma^0 u(\mathbf{p}, s)$ while $v(-\mathbf{p}, s) = -\gamma^0 v(\mathbf{p}, s)$.
- (c) Now show that eq. (15) implies

$$\begin{aligned} \widehat{\mathbf{P}} \hat{a}_{\mathbf{p},s} \widehat{\mathbf{P}} &= \pm \hat{a}_{-\mathbf{p},+s}, & \widehat{\mathbf{P}} \hat{a}_{\mathbf{p},s}^\dagger \widehat{\mathbf{P}} &= \pm \hat{a}_{-\mathbf{p},+s}^\dagger, \\ \widehat{\mathbf{P}} \hat{b}_{\mathbf{p},s} \widehat{\mathbf{P}} &= \mp \hat{b}_{-\mathbf{p},+s}, & \widehat{\mathbf{P}} \hat{b}_{\mathbf{p},s}^\dagger \widehat{\mathbf{P}} &= \mp \hat{b}_{-\mathbf{p},+s}^\dagger, \end{aligned} \quad (16)$$

and hence

$$\widehat{\mathbf{P}} |F(\mathbf{p}, s)\rangle = \pm |F(-\mathbf{p}, +s)\rangle \quad \text{and} \quad \widehat{\mathbf{P}} |\overline{F}(\mathbf{p}, s)\rangle = \mp |\overline{F}(-\mathbf{p}, +s)\rangle. \quad (17)$$

Note that the fermion F and the antifermion \overline{F} have opposite intrinsic parities!

5. A Dirac spinor field $\Psi(x)$ comprises two 2-component Weyl spinor fields,

$$\hat{\Psi}(x) = \begin{pmatrix} \hat{\psi}_L(x) \\ \hat{\psi}_R(x) \end{pmatrix}. \quad (18)$$

Spell out the actions of \mathbf{C} , \mathbf{P} , and \mathbf{CP} (combined —bf *CandP* symmetries on the Weyl spinors. In particular, show that \mathbf{C} and \mathbf{P} interchange the two spinors, while the combined \mathbf{CP} symmetry acts on the ψ_L and the ψ_R independently from each other.

6. Consider the bilinear products of a Dirac field $\Psi(x)$ and its conjugate $\bar{\Psi}(x)$. Generally, such products have form $\bar{\Psi}\Gamma\Psi$ where Γ is one of 16 matrices discussed in the [previous homework#7](#) (problem 4.g); altogether, we have

$$S = \bar{\Psi}\Psi, \quad V^\mu = \bar{\Psi}\gamma^\mu\Psi, \quad T^{\mu\nu} = \bar{\Psi}\frac{i}{2}\gamma^{[\mu}\gamma^{\nu]}\Psi, \quad A^\mu = \bar{\Psi}\gamma^5\gamma^\mu\Psi, \quad P = \bar{\Psi}i\gamma^5\Psi. \quad (19)$$

(a) Show that all the bilinears (19) are Hermitian.

Hint: First, show that $(\bar{\Psi}\Gamma\Psi)^\dagger = \bar{\Psi}\Gamma\Psi$.

Note: despite the Fermi statistics, $(\Psi_\alpha^\dagger\Psi_\beta)^\dagger = +\Psi_\beta^\dagger\Psi_\alpha$.

(b) Show that under *continuous* Lorentz symmetries, the S and the P transform as scalars, the V^μ and the A^μ as vectors, and the $T^{\mu\nu}$ as an antisymmetric tensor.

(c) Find the transformation rules of the bilinears (19) under parity and show that while S is a true scalar and V is a true (polar) vector, P is a pseudoscalar and A is an axial vector.

Finally, consider the charge-conjugation properties of the Dirac bilinears. To avoid the operator-ordering problems, take the classical limit where $\Psi(x)$ and $\Psi^\dagger(x)$ *anticommute* with each other, $\Psi_\alpha\Psi_\beta^\dagger = -\Psi_\beta^\dagger\Psi_\alpha$.

(d) Show that \mathbf{C} turns $\bar{\Psi}\Gamma\Psi$ into $\bar{\Psi}\Gamma^c\Psi$ where $\Gamma^c = \gamma^0\gamma^2\Gamma^\top\gamma^0\gamma^2$.

(e) Calculate Γ^c for all 16 independent matrices Γ and find out which Dirac bilinears are \mathbf{C} -even and which are \mathbf{C} -odd.