

1. In class we have focused on QCD and QCD-like theories of non-abelian gauge fields coupled to Dirac fermions in some multiplet(s) of the gauge group G , cf. [my notes on QCD Feynman rules and Ward identities](#). This problem is about the scalar QCD, or more generally a non-abelian gauge theory with some gauge group G and complex scalar fields $\Phi^i(x)$ in some multiplet (r) of G .

- (a) Write down the physical Lagrangian of this theory, the complete bare Lagrangian of the quantum theory in the Feynman gauge, and the Feynman rules.

Now consider the annihilation process $\Phi + \Phi^* \rightarrow 2$ gauge bosons. At the tree level, there are four Feynman diagrams contributing to this process.

- (b) Draw the diagrams and write down the tree-level annihilation amplitude.

As discussed in class, amplitudes involving the non-abelian gauge fields satisfy a weak form of the Ward identity: *On-shell Amplitudes involving a longitudinally polarized gauge bosons vanish, provided all the other gauge bosons are transversely polarized*. In other words,

$$\mathcal{M} \equiv e_1^{\mu_1} e_2^{\mu_2} \cdots e_n^{\mu_n} \mathcal{M}_{\mu_1 \mu_2 \cdots \mu_n}(\text{momenta}) = 0$$

when $e_1^\mu \propto k_1^\mu$ but $e_2^\nu k_{2\nu} = \cdots = e_n^\nu k_{n\nu} = 0$.

- (c) Verify this identity for the scalar annihilation amplitude: Show that IF $e_2^\nu k_{2\nu} = 0$ THEN $k_{1\mu} \mathcal{M}^{\mu\nu} e_{2\nu} = 0$.

Similar to what we had in class for the quark-antiquark annihilations, there are non-zero amplitudes for the scalar ‘quark’ and ‘antiquark’ annihilating into a pair of longitudinal gluons or a ghost-antighost pair, but the crosssections for these two unphysical processes cancel each other.

- (d) Take both final-state gluons to be longitudinally polarized; specifically, assume null polarization vectors $e_1^\mu = (1, +\mathbf{n}_1)/\sqrt{2}$ for the first gluon and $e_2^\nu = (1, -\mathbf{n}_2)/\sqrt{2}$ for the second gluons.

Calculate the tree-level annihilation amplitude $\Phi + \Phi^* \rightarrow g_L + g_L$ for these polarizations.

- (e) Next, calculate the tree amplitude for the $\Phi + \Phi^* \rightarrow \text{gh} + \overline{\text{gh}}$. There is only one tree graph for this process, so evaluating it should not be hard.
- (f) Compare the two un-physical amplitudes and show that the corresponding partial cross-sections cancel each other, thus

$$\frac{d\sigma_{\text{net}}}{d\Omega} = \frac{d\sigma_{\text{physical}}}{d\Omega}. \quad (1)$$

2. Next, an exercise in group theory you would need for QCD and QCD-like gauge theories. Consider a generic simple non-abelian compact Lie group G and its generators T^a . For a suitable normalization of the generators,

$$\text{tr}_{(r)}(T^a T^b) \equiv \text{tr} \left(T_{(r)}^a T_{(r)}^b \right) = R(r) \delta^{ab} \quad (2)$$

where the trace is taken over any complete multiplet (r) — irreducible or reducible, it does not matter — and $T_{(r)}^a$ is the matrix representing the generator T^a in that multiplet. The coefficient $R(r)$ in eq. (2) depends on the multiplet (r) but it's the same for all generators T^a and T^b . The $R(r)$ is called the *index* of the multiplet (r) .

The (quadratic) Casimir operator $C_2 = \sum_a T^a T^a$ commutes with all the generators, $\forall b, [C_2, T^b] = 0$. Consequently, when we restrict this operator to any *irreducible* multiplet (r) of the group G , it becomes a unit matrix times some number $C(r)$. In other words,

$$\text{for an irreducible } (r), \quad \sum_a T_{(r)}^a T_{(r)}^a = C(r) \times \mathbf{1}_{(r)}. \quad (3)$$

For example, for the isospin group $SU(2)$, the Casimir operator is $C_2 = \vec{I}^2$, the irreducible multiplets have definite isospin $I = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \dots$, and $C(I) = I(I + 1)$.

- (a) Show that for any irreducible multiplet (r) ,

$$\frac{R(r)}{C(r)} = \frac{\dim(r)}{\dim(G)}. \quad (4)$$

In particular, for the $SU(2)$ group, this formula gives $R(I) = \frac{1}{3}I(I + 1)(2I + 1)$.

- (b) Suppose the first three generators T^1 , T^2 , and T^3 of G generate an $SU(2)$ subgroup, thus

$$[T^1, T^2] = iT^3, \quad [T^2, T^3] = iT^1, \quad [T^3, T^1] = iT^2. \quad (5)$$

Show that if a multiplet (r) of G decomposes into several $SU(2)$ multiplets of isospins I_1, I_2, \dots, I_n , then

$$R(r) = \sum_{i=1}^n \frac{1}{3} I_i(I_i + 1)(2I_i + 1). \quad (6)$$

- (c) Now consider the $SU(N)$ group with an obvious $SU(2)$ subgroup of matrices acting only on the first two components of a complex N -vector. This complex N -vector is called the fundamental multiplet (of the $SU(N)$) and denoted (N) or \mathbf{N} . As far as the $SU(2)$ subgroup is concerned, (N) comprises one doublet and $N - 2$ singlets, hence

$$R(N) = \frac{1}{2} \quad \text{and} \quad C(N) = \frac{N^2 - 1}{2N}. \quad (7)$$

Show that the adjoint multiplet of the $SU(N)$ decomposes into one $SU(2)$ triplet, $2(N - 2)$ doublets, and $(N - 2)^2$ singlets, therefore

$$R(\text{adj}) = C(\text{adj}) \equiv C(G) = N. \quad (8)$$

Hint: $(N) \times (\overline{N}) = (\text{adj}) + (1)$.

- (d) The symmetric and the anti-symmetric 2-index tensors form irreducible multiplets of the $SU(N)$ group. Find out the decomposition of these multiplets under the $SU(2) \subset SU(N)$ and calculate their respective indices R and Casimirs C .
3. Now let's apply this group theory to physics. Consider quark-antiquark pair production in QCD, specifically $u\bar{u} \rightarrow d\bar{d}$. There is only one tree diagram contributing to this process,



Evaluate this diagram, then sum/average the $|\mathcal{M}|^2$ over both spins and *colors* of the final/initial particles to calculate the total cross section. For simplicity, you may neglect the

quark masses.

Note that the diagram (9) looks exactly like the QED pair production process $e^-e^+ \rightarrow$ virtual $\gamma \rightarrow \mu^-\mu^+$, so you can re-use the QED formula for summing/averaging over the spins, *cf.* [my notes on Dirac traceology from the Fall semester](#), pages 10–13. But in QCD, you should also sum/average over the colors of all the quarks, and that's the whole point of this exercise.