


This homework is about one-dimensional square wells and barriers. For most students, this should be a refresher of undergraduate quantum mechanics.

1. Consider the bound states in an asymmetric square well,



A symmetric potential well in one dimension always has at least one bound state, but this is not always true for the asymmetric wells with $V_1 \neq V_2$.

- (a) Solve the Schrödinger equation in each region of x , apply the ‘boundary’ conditions at $x \rightarrow \pm\infty$, and spell out the continuity conditions at $x = 0$ and $x = L$. Assume a bound state, thus $E < V_1 \leq V_2$.
- (b) Solve the continuity conditions and show that the bound states energies — and the corresponding k , κ_1 , and κ_2 — must obey


$$kL = \arctan \frac{\kappa_1}{k} + \arctan \frac{\kappa_2}{k} + n \times \pi = \arccos \sqrt{\frac{E}{V_1}} + \arccos \sqrt{\frac{E}{V_2}} + n \times \pi \quad (2)$$

for an integer n

The best way to solve eq. (2) is graphic: plot both sides of the equation as functions of k — or as functions of $\sqrt{E/V_1} = k \times \text{const}$ — and look for the intersections. Note that the RHS has a separate branch for each n , so there could be several intersections and hence several bound states with different energies.

- (c) Show that for a symmetric well with $V_2 = V_1$ and any $L > 0$, the branch with $n = 0$ always has an intersection. Thus, a symmetric well always has at least one bound state.
- (d) On the other hand, an asymmetric well with $V_1 \neq V_2$ and small enough L has no bound state at all. Show this, and also calculate the smallest width L_{\min} of the asymmetric well that does have a bound state.

2. Next, consider a symmetric square barrier,



$$V(x) = \begin{cases} V_b & \text{for } 0 < x < L, \\ 0 & \text{for } x < 0 \text{ or } x > L. \end{cases}$$

(3)

Calculate the reflection and transmission coefficients for this barrier as functions of the unbound state energy $E > 0$ for three cases:

- (a) $E < V_b$, tunneling under the barrier.
- (b) $E > V_b$, flying over the barrier.
- (c) The barrier is actually a well with $V_b < 0$.

Hint: show that in terms of k' (inside the barrier/well) and k (outside), the wave function and the continuity equations are exactly in part (b), then re-use the result of part (b) instead of re-calculating it from scratch.

★ Also, show that the reflection and transmission coefficients as functions of energy in all 3 cases are related by analytic continuation.