

# OPTICAL THEOREM

The optical theorem relates the imaginary part of the elastic scattering amplitude in the forward direction to the total cross-section — elastic + inelastic — of the initial particles. In the non-relativistic normalization,

$$\text{Im } f_{\text{elastic}}(\theta = 0) = \frac{k_{\text{reduced}}}{4\pi} \times \sigma_{\text{total}}. \quad (1)$$

You have probably seen this formula in an undergraduate QM class in the context of the partial wave analysis of potential scattering; but in case you have never studied the subject, here are [my notes on the partial wave analysis](#) from a graduate E&M class (387 K) I have recently taught. Anyway, for a purely elastic scattering

$$f(\theta) = \frac{1}{k} \sum_{\ell=0}^{\infty} (2\ell + 1) P_{\ell}(\cos \theta) \times \left( \frac{e^{2i\delta_{\ell}} - 1}{2i} = e^{i\delta_{\ell}} \sin \delta_{\ell} \right), \quad (2)$$

where  $\delta_{\ell}$  is the phase shift for the partial wave  $\ell$ . Note that for small phase shifts  $\delta_{\ell} \ll 1$ , the real part of the scattering amplitude is much larger than its imaginary part,  $\text{Re } f \sim \delta$  while  $\text{Im } f \sim \delta^2$ . For the scattering amplitude (2), the total cross-section is

$$\sigma_{\text{total}} = \int |f|^2 d^2\Omega = \frac{4\pi}{k^2} \times \sum_{\ell=0}^{\infty} (2\ell + 1) \sin^2 \delta_{\ell}, \quad (3)$$

while the imaginary part of the forward scattering amplitude is

$$\text{Im } f(\theta = 0) = \frac{1}{k} \times \sum_{\ell=0}^{\infty} (2\ell + 1) \sin^2 \delta_{\ell}, \quad (4)$$

in manifest agreement with the optical theorem (1).

In relativistic notations, the optical theorem (1) becomes

$$\text{Im } \mathcal{M} \left( \begin{array}{c} \text{elastic} \\ \text{forward} \end{array} \right) = 2E_1 E_2 |\mathbf{v}_1 - \mathbf{v}_2| \times \sigma_{\text{total}}. \quad (5)$$

where the pre-factor  $2E_1 E_2 |\mathbf{v}_1 - \mathbf{v}_2|$  is invariant under Lorentz boosts along the axis of collision. In the center-of-mass frame, it amounts to  $4E_{\text{cm}}^{\text{net}} |\mathbf{p}_{\text{cm}}|$ .

## Proof

The optical theorem follows from the unitarity of the S-matrix — or rather, the scattering operator, —  $\hat{S}^\dagger \hat{S} = 1$ . To see how this works, let's separate the scattering from non-scattering events,  $\hat{S} = 1 + i\hat{T}$ , and express the unitarity of  $\hat{S}$  in terms of  $\hat{T}$ :

$$1 = (\hat{S}^\dagger = 1 - \hat{T}^\dagger)(\hat{S} = 1 + i\hat{T}) = 1 + i\hat{T} - i\hat{T}^\dagger + \hat{T}^\dagger \hat{T}, \quad (6)$$

hence

$$i\hat{T}^\dagger - i\hat{T} = \hat{T}^\dagger \hat{T}. \quad (7)$$

Now let's take the diagonal matrix elements  $\langle i | \dots | i \rangle$  of both sides of this formula for some state  $|i\rangle$ , which we shall eventually take to be the initial state  $|1 + 2\rangle$  of two particles about to collide. On the LHS of eq. (7)

$$\langle i | i\hat{T}^\dagger - i\hat{T} | i \rangle = i \langle i | \hat{T} | i \rangle^* - i \langle i | \hat{T} | i \rangle = 2 \text{Im} \langle i | \hat{T} | i \rangle \quad (8)$$

while on the RHS of eq. (7)

$$\langle i | \hat{T}^\dagger \hat{T} | i \rangle = \sum_{|f\rangle} \langle i | \hat{T}^\dagger | f \rangle \langle f | \hat{T} | i \rangle = \sum_{|f\rangle} \left| \langle f | \hat{T} | i \rangle \right|^2, \quad (9)$$

hence

$$2 \text{Im} \langle i | \hat{T} | i \rangle = \sum_{|f\rangle} \left| \langle f | \hat{T} | i \rangle \right|^2. \quad (10)$$

Physically, the sum on the RHS here is over all possible final states of the initial particles' scattering.

Next, let's factor out the energy-momentum conservation from the matrix elements of the  $\hat{T}$  operator,

$$\langle f | \hat{T} | i \rangle = (2\pi)^4 \delta^{(4)}(p_f - p_i) \times \langle f | \widehat{\mathcal{M}} | i \rangle. \quad (11)$$

Plugging this formula directly into eq. (10) gives us

$$2 \text{Im} \langle i | \widehat{\mathcal{M}} | i \rangle \times (2\pi)^4 \delta^{(4)}(p_i - p_i) = \sum_{|f\rangle} \left| \langle f | \widehat{\mathcal{M}} | i \rangle \right|^2 \times \left( (2\pi)^4 \delta^{(4)}(p_f - p_i) \right)^2, \quad (12)$$

with troublesome  $\delta$ -functions on both sides of the equation. To resolve this trouble, consider

the state  $|i'\rangle$  which is very similar to the initial state  $|i\rangle$  except for a possible tiny difference in the net momentum  $p_{i'}$ . By similar I mean similar enough that we may approximate

$$\langle i' | \widehat{\mathcal{M}} | i \rangle = \langle i | \widehat{\mathcal{M}} | i \rangle \quad \text{and} \quad \langle i' | \widehat{\mathcal{M}} | f \rangle = \langle i | \widehat{\mathcal{M}} | f \rangle \quad \forall |f\rangle. \quad (13)$$

Consequently, taking the slightly-off-diagonal matrix elements  $\langle i' | \dots | i \rangle$  of both sides of eq. (7), we get on the LHS

$$\langle i' | i\hat{T}^\dagger - i\hat{T} | i \rangle = (2\pi)^4 \delta^{(4)}(p_{i'} - p_i) \times \langle i | i\widehat{\mathcal{M}}^\dagger - i\widehat{\mathcal{M}} | i \rangle = (2\pi)^4 \delta^{(4)}(p_{i'} - p_i) \times 2 \text{Im} \langle i | \widehat{\mathcal{M}} | i \rangle, \quad (14)$$

and on the RHS

$$\begin{aligned} \langle i' | \hat{T}^\dagger \hat{T} | i \rangle &= \sum_{|f\rangle} \langle f | \hat{T} | i' \rangle^* \langle f | \hat{T} | i \rangle \\ &= \sum_{|f\rangle} \left| \langle f | \widehat{\mathcal{M}} | i \rangle \right|^2 \times (2\pi)^4 \delta^{(4)}(p_f - p_{i'}) \times (2\pi)^4 \delta^{(4)}(p_f - p_i) \\ &= (2\pi)^4 \delta^{(4)}(p_{i'} - p_i) \times \sum_{|f\rangle} \left| \langle f | \widehat{\mathcal{M}} | i \rangle \right|^2 \times (2\pi)^4 \delta^{(4)}(p_f - p_i). \end{aligned} \quad (15)$$

Equating the last two formulae and dropping the overall factor  $(2\pi)^4 \delta^{(4)}(p_{i'} - p_i)$  from both, we arrive at the improved version of eq. (12), namely

$$2 \text{Im} \langle i | \widehat{\mathcal{M}} | i \rangle = \sum_{|f\rangle} \left| \langle f | \widehat{\mathcal{M}} | i \rangle \right|^2 \times (2\pi)^4 \delta^{(4)}(p_f - p_i). \quad (16)$$

Now let  $|i\rangle = |1 + 2\rangle$  be the initial state of two particles about to collide, so that  $|f\rangle$  runs over all possible final states of this collision. Thus, summing over  $|f\rangle$  means summing over all possible reaction *channels* — *i.e.*, sets of final particles, — and then for each final-state particle integrating over its momentum with the relativistic measure and summing over its

discrete quantum numbers like spin:

$$\begin{aligned} \langle f | &= \langle \text{channel} : (p', s')_1, \dots, (p', s')_n |, \\ \sum_{\langle f |} &= \sum_{\text{channels}} \prod_{a=1}^n \left( \sum_{s'_a} \int \frac{d^3 \mathbf{p}'_a}{(2\pi)^3 2E'_a} \right). \end{aligned} \quad (17)$$

Consequently, eq. (16) becomes

$$\begin{aligned} 2 \text{Im} \langle i | \widehat{\mathcal{M}} | i \rangle &= \sum_{\text{channels}} \prod_{a=1}^n \left( \sum_{s'_a} \int \frac{d^3 \mathbf{p}'_a}{(2\pi)^3 2E'_a} \right) \left| \langle \text{channel} : (p', s')_1, \dots, (p', s')_n | \widehat{\mathcal{M}} | i \rangle \right|^2 \times \\ &\quad \times (2\pi)^4 \delta^{(4)}(p'_1 + \dots + p'_n - p_i^{\text{net}}). \end{aligned} \quad (18)$$

Note that the momentum integral (and spin sum) on the RHS here is precisely the phase space integral (and spin sum) for the scattering process  $(1 + 2 \rightarrow 1' + \dots + n')$ . Specifically,

$$\begin{aligned} \sigma_{\text{net}}(1 + 2 \rightarrow 1' + \dots + n') &= \frac{1}{4E_1 E_2 |\mathbf{v}_{12}^{\text{rel}}|} \prod_{a=1}^n \left( \sum_{s'_a} \int \frac{d^3 \mathbf{p}'_a}{(2\pi)^3 2E'_a} \right) \left| \langle (p', s')_1, \dots, (p', s')_n | \widehat{\mathcal{M}} | 1 + 2 \rangle \right|^2 \times \\ &\quad \times (2\pi)^4 \delta^{(4)}(p'_1 + \dots + p'_n - p_i^{\text{net}}), \end{aligned} \quad (19)$$

*cf.* [my notes on the phase space](#). Therefore, we may rephrase eq. (18) in terms of the net cross-sections for each channel and ultimately in terms of the total cross-section for all the channel combined:

$$\begin{aligned} 2 \text{Im} \langle 1 + 2 | \widehat{\mathcal{M}} | 1 + 2 \rangle &= \sum_{\text{channels}} 4E_1 E_2 |\mathbf{v}_{12}^{\text{rel}}| \times \sigma_{\text{net}}(1 + 2 \rightarrow 1' + \dots + n') \\ &= 4E_1 E_2 |\mathbf{v}_{12}^{\text{rel}}| \times \sum_{\text{channels}} \sigma_{\text{net}}(1 + 2 \rightarrow 1' + \dots + n') \\ &= 4E_1 E_2 |\mathbf{v}_{12}^{\text{rel}}| \times \sigma_{\text{total}}(1 + 2 \rightarrow \text{anything}). \end{aligned} \quad (20)$$

Finally, we identify the matrix element on the LHS here as the amplitude for the forward elastic scattering  $|1 + 2\rangle \rightarrow |\text{exactly the same state}\rangle$ , thus

$$\text{Im} \mathcal{M} \begin{pmatrix} \text{elastic} \\ \text{forward} \end{pmatrix} = 2E_1 E_2 |\mathbf{v}_{12}^{\text{rel}}| \times \sigma_{\text{total}}(1 + 2 \rightarrow \text{anything}). \quad (21)$$

And this completes my proof of the optical theorem.

BTW, there is a similar version of the optical theorem for the decays of unstable particles. Indeed, let in eq. (18)  $|i\rangle$  be the initial state of a single unstable particle (instead of two particles about to collide). Then the RHS of eq. (18) becomes the phase space integral (and the spin sum) for the net decay rate of the initial particle into all the available decay channels. Indeed,

$$\Gamma_{\text{net}}(1 \rightarrow 1' + \dots + n') = \frac{1}{2M} \prod_{a=1}^n \left( \sum_{s'_a} \int \frac{d^3 \mathbf{p}'_a}{(2\pi)^3 2E'_a} \right) \left| \langle (p', s')_1, \dots, (p', s')_n | \widehat{\mathcal{M}} | 1 \rangle \right|^2 \times \\ \times (2\pi)^4 \delta^{(4)}(p'_1 + \dots + p'_n - p_i^{\text{net}}), \quad (22)$$

— *cf. my notes on the phase space*, — hence

$$\begin{aligned} \text{Im} \langle 1 | \widehat{\mathcal{M}} | 1 \rangle &= \frac{1}{2} \sum_{\text{channels}} 2M \times \Gamma_{\text{net}}(1 \rightarrow 1' + \dots + n') \\ &= M \times \sum_{\text{channels}} \Gamma_{\text{net}}(1 \rightarrow 1' + \dots + n') \\ &= M \times \Gamma_{\text{total}}(1 \rightarrow \text{anything}). \end{aligned} \quad (23)$$

We shall return to this formula in a few lectures, once we have a better understanding of the one-particle to one-particle amplitudes.

## Application to the $\lambda\phi^4$ Theory.

In potential scattering, the first Born approximation yields a real scattering amplitude, but at the higher orders of perturbation theory the amplitude becomes complex, with both real and imaginary parts. Likewise, in quantum field theory the tree-level amplitudes are real, — at least for the elastic scattering in the forward direction, — but the loop corrections make for complex amplitudes. The reason for this behavior is the optical theorem (21), plus power-of-the-coupling counting in the perturbation theory. Let's see how this works for the  $\lambda\phi^4$  theory.

For the elastic scattering  $\mathcal{M}_{\text{tree}} = -\lambda$  (regardless of momenta) while the loop corrections are of the higher order in  $\lambda$ , hence

$$\frac{d\sigma^{\text{elastic}}}{d\Omega_{\text{cm}}} = \frac{\lambda^2 + O(\lambda^3)}{64\pi^2 s} \quad (24)$$

and

$$\sigma_{\text{net}}^{\text{elastic}} = \frac{\lambda^2 + O(\lambda^3)}{64\pi^2 s} \times \frac{4\pi}{2} = \frac{\lambda^2 + O(\lambda^3)}{32\pi s}. \quad (25)$$

For the inelastic processes  $2 \rightarrow n$  ( $n \geq 4$ ), the tree amplitude is  $O(\lambda^{n/2})$  and the loop corrections are of higher order, hence  $\mathcal{M} = O(\lambda^{n/2})$  and

$$\sigma^{\text{inelastic}} = O(\lambda^n) \quad \text{for } n \geq 4. \quad (26)$$

Consequently,

$$\sigma_{\text{total}} = \sigma_{\text{net}}^{\text{elastic}} + O(\lambda^4) = \frac{\lambda^2 + O(\lambda^3)}{32\pi s}. \quad (27)$$

By the optical theorem, this gives us the imaginary part of the forward elastic amplitude as

$$\text{Im } \mathcal{M} \left( \begin{array}{c} \text{elastic} \\ \text{forward} \end{array} \right) = 2E_1 E_2 |\mathbf{v}_{12}^{\text{rel}}| \times \frac{\lambda^2 + O(\lambda^3)}{32\pi s}, \quad (28)$$

where in the center of mass frame

$$2E_1 E_2 |\mathbf{v}_{12}^{\text{rel}}| = 2E^2 \times 2v = s \times v \quad (29)$$

(where  $v$  is the speed of each particle relative to the CM), hence

$$\text{Im } \mathcal{M} \left( \begin{array}{c} \text{elastic} \\ \text{forward} \end{array} \right) = \frac{\lambda^2 v}{32\pi} + O(\lambda^3). \quad (30)$$

In terms of the loop counting for the amplitude on the LHS here, this means

$$\text{Im } \mathcal{M}_{\text{tree}} \left( \begin{array}{c} \text{elastic} \\ \text{forward} \end{array} \right) = 0, \quad (31)$$

$$\text{while } \text{Im } \mathcal{M}_{1\text{loop}} \left( \begin{array}{c} \text{elastic} \\ \text{forward} \end{array} \right) = \frac{\lambda^2 v}{32\pi} > 0. \quad (32)$$

Note: the optical theorem not only explains why the tree amplitude is real while the one-loop amplitude is complex, it also gives us specific predictions for the imaginary part of the one-loop amplitude in terms of the tree-level total cross-section. In the same way, it would give us a specific prediction for the imaginary part of the two-loop amplitude in terms of the one-loop total cross-section, *etc.*, *etc.* But let's not get too far into the higher-order calculations.

Instead, let's verify the tree-level and 1-loop level predictions (31) and (32). At the tree level, we indeed have

$$\mathcal{M}_{\text{tree}}^{\text{elastic}} = -\lambda, \quad \text{Im } \mathcal{M}_{\text{tree}}^{\text{elastic}} = 0 \quad (33)$$

for any scattering angle  $\theta$  and not just  $\theta = 0$ . Likewise, we shall see in a moment that

$$\forall \theta : \quad \text{Im } \mathcal{M}_{1\text{loop}}^{\text{elastic}} = \frac{\lambda^2 v}{32\pi} \quad (34)$$

where

$$v = \frac{|\mathbf{p}|}{E} = \sqrt{1 - \frac{m^2}{E^2}} = \sqrt{1 - \frac{4m^2}{s}}. \quad (35)$$

Indeed, we have seen earlier in class that

$$\mathcal{M}_{1\text{loop}}^{\text{elastic}} = \frac{\lambda^2}{32\pi^2} \left( J(4) - J(t/m^2) - J(u/m^2) - J(s/m^2) \right), \quad (36)$$

where

$$J(t/m^2) = \int_0^1 dx \log \frac{m^2 - tx(1-x)}{m^2} \quad (37)$$


and likewise for  $J(u/m^2)$  and  $J(s/m^2)$ . For elastic scattering  $t < 0$ , hence

$$J(t) = \int_0^1 dx \log \frac{m^2 + \text{positive}}{m^2}, \quad (38)$$

which is real and positive. Likewise,  $u < 0$  and hence  $J(u/m^2)$  is real and positive. But  $s > +4m^2$ , which makes  $J(s/m^2)$  complex rather than real. Indeed,

$$J(s/m^2) = \int_0^1 dx \log \frac{m^2 - sx(1-x)}{m^2}, \quad (39)$$

and for  $s > +4m^2$  the argument of the logarithm here becomes negative for some  $x$ :




$$x_{1,2} = \frac{1}{2} \mp \sqrt{\frac{1}{4} - \frac{m^2}{s}} = \frac{1 \mp v}{2}. \quad (40)$$

For a negative or complex argument the logarithm becomes complex,

$$\log(z) = \log(|z|) + i \arg(z), \quad (41)$$

which has a branch cut in the complex  $z$  plane along the negative half of the real axis,



thus

$$\log(\text{negative} \pm i\epsilon) = \text{real} \pm i\pi. \quad (42)$$

Consequently,

$$\text{Im} \log \frac{m^2 - sx(1-x)}{m^2} = \begin{cases} \pm\pi & \text{for } x_1 < x < x_2, \\ 0 & \text{otherwise,} \end{cases} \quad (43)$$

and therefore

$$\text{Im} J(s/m^2) = \int_0^1 dx \text{Im} \log \frac{m^2 - sx(1-x)}{m^2} = \pm\pi \times (x_2 - x_1) = \pm\pi v \quad (44)$$

(where  $v$  is the particles' speed in the CM frame). The sign of this imaginary part depends on the side of the logarithm's branch cut we end up on for negative  $m^2 - sx(1-x)$ , which



in turn depends on the analytic continuation from a real  $s$  to  $s + i\epsilon$  or to  $s - i\epsilon$ :

$$\begin{aligned} \text{for } s \rightarrow s + i\epsilon, \quad m^2 - sx(1-x) = \text{real} - i\epsilon &\implies \log = \text{real} - i\pi, \\ \text{while for } s \rightarrow s - i\epsilon, \quad m^2 - sx(1-x) = \text{real} + i\epsilon &\implies \log = \text{real} + i\pi, \end{aligned} \quad (45)$$

and hence

$$\text{Im } J((s \pm i\epsilon)/m^2) = \mp \pi v. \quad (46)$$

In the context of the one-loop elastic amplitude (36), this means

$$\text{Im } \mathcal{M}_{1\text{loop}}^{\text{elastic}}(s \pm i\epsilon, t) = -\frac{\lambda^2}{32\pi^2} \text{Im } J((s \pm i\epsilon)/m^2) = \pm \frac{\lambda^2 v}{32\pi}. \quad (47)$$

As we see, this formula is in perfect agreement with the optical theorem's prediction (32), *provided we interpret  $s$  as  $s + i\epsilon$ .*

The reason for the choice of  $s \rightarrow s + i\epsilon$  rather than  $s \rightarrow s - i\epsilon$  stems from the origin of the  $J(s/m^2)$  in the  $s$ -channel amplitude

$$\mathcal{F}(s) = \frac{\lambda^2}{2} \int_0^1 dx \int_{\text{reg}} \frac{d^4 k}{(2\pi)^4} \frac{-i}{[k^2 - \Delta(x, s) + i\epsilon]^2}, \quad (48)$$

*cf.* eq. (20) of [my introductions to the one-loop amplitudes](#) for a similar  $t$ -channel amplitude. After the Wick rotation to the Euclidean momentum space, eq. (48) becomes

$$\mathcal{F}(s) = \frac{\lambda^2}{2} \int_0^1 dx \int_{\text{reg}} \frac{d^4 k_E}{(2\pi)^4} \frac{1}{[k_E^2 + \Delta(x, s) - i\epsilon]^2}. \quad (49)$$

For a similar  $t$ -channel amplitude, we may disregard the  $-i\epsilon$  term in the denominator because  $\Delta(x, t) > 0$  for all  $x$  so there are no poles in the Euclidean momentum space. But in the  $s$  channel  $\Delta(x, s)$  turns negative for some  $x$ , so we should retain the  $-i\epsilon$  term. Hence, in all

the following formulae for the  $s$  channel we should replace  $\Delta(x, s)$  with

$$\Delta(x, s) - i\epsilon = m^2 - x(x-x) \times s - i\epsilon = m^2 - x(1-x) \times (s + i\epsilon), \quad (50)$$

which is equivalent to replacing  $s \rightarrow s + i\epsilon$ . And that's how we end up with  $J((s + i\epsilon)/m^2)$  in the  $s$ -channel amplitude and therefore

$$\mathcal{M}_{1\text{loop}}^{\text{elastic}} = \frac{\lambda^2}{32\pi^2} \left( J(4) - J(t/m^2) - J(u/m^2) - J((s + i\epsilon)/m^2) \right). \quad (51)$$

In my [next set of notes about the field correlation functions](#) I shall give you a more general reason for always interpreting  $s$  as  $s + i\epsilon$  rather than  $s - \epsilon$ .