

1. In the [previous homework](#) — set#5, problem#4(c) — we saw that the little group of Lorentz symmetries preserving the lightlike momentum p^μ or a massless particle is generated by the 3 components of the vector

$$\hat{\mathbf{I}} = \hat{\mathbf{J}} + \vec{\beta} \times \hat{\mathbf{K}}. \quad (1)$$

In particular, the component in the direction of the particle's velocity

$$\hat{I}_{\parallel} = \vec{\beta} \cdot \hat{\mathbf{I}} = \vec{\beta} \cdot \hat{\mathbf{J}} = \hat{\lambda} \quad (2)$$

is the helicity operator. As I explained in class, the finite unitary representations of this little group are singlets of definite helicity, specifically the states $|p, \lambda\rangle$ obeying

$$\hat{I}_{\parallel} |p, \lambda\rangle = \lambda |p, \lambda\rangle \quad \text{and} \quad \hat{\mathbf{I}}_{\perp} |p, \lambda\rangle = 0. \quad (3)$$

- (a) Show that in 4D terms, the conditions (3) amount to

$$\frac{1}{2} \epsilon_{\mu\alpha\beta\gamma} \hat{J}^{\alpha\beta} \hat{P}^{\gamma} |p, \lambda\rangle = \lambda \hat{P}_{\mu} |p, \lambda\rangle. \quad (4)$$

- (b) Use eq. (4) to show that the *continuous* Lorentz transforms do not change helicities of *massless* particles,

$$\forall L \in \text{SO}^+(3, 1), \quad \hat{\mathcal{D}}(L) |p, \lambda\rangle = |Lp, \text{same } \lambda\rangle \times e^{i \text{phase}}. \quad (5)$$

2. The Spin(3, 1) group — the double cover of the continuous Lorentz group $SO^+(3, 1)$ — is isomorphic to $SL(2, \mathbf{C})$, the group of complex (but not necessarily unitary) 2×2 matrices of unit determinant. The relations between such matrices and the Lorentz symmetries are explained in [my notes on Lorentz representations for the fields](#), but some technical details were left out as exercises for the students. This problem collects these exercises.

(a) Show that the components of the two 3-vectors

$$\hat{\mathbf{J}}_+ = \frac{1}{2}(\hat{\mathbf{J}} + i\hat{\mathbf{K}}) \quad \text{and} \quad \hat{\mathbf{J}}_- = \frac{1}{2}(\hat{\mathbf{J}} - i\hat{\mathbf{K}}) = \hat{\mathbf{J}}_+^\dagger. \quad (6)$$

obey commutation relations

$$[\hat{J}_+^i, \hat{J}_+^j] = i\epsilon^{ijk} \hat{J}_+^k, \quad [\hat{J}_-^i, \hat{J}_-^j] = i\epsilon^{ijk} \hat{J}_-^k, \quad \text{but} \quad [\hat{J}_+^i, \hat{J}_-^j] = 0. \quad (7)$$

Now consider the $\mathbf{2}$ ($j_+ = \frac{1}{2}, j_- = 0$) and the $\bar{\mathbf{2}}$ ($j_+ = 0, j_- = \frac{1}{2}$) representations of the Lorentz or rather Spin(3, 1) group. In the $\mathbf{2}$ representation $\mathbf{J} = \frac{1}{2}\boldsymbol{\sigma}$ and $\mathbf{K} = -\frac{i}{2}\boldsymbol{\sigma}$ while in the $\bar{\mathbf{2}}$ representation $\mathbf{J} = \frac{1}{2}\boldsymbol{\sigma}$ and $\mathbf{K} = +\frac{i}{2}\boldsymbol{\sigma}$, hence for a 3-space rotation R through angle ϕ around axis \mathbf{n}

$$M_{\mathbf{2}}(R) = M_{\bar{\mathbf{2}}} = \exp\left(-\frac{i}{2}\phi\mathbf{n} \cdot \boldsymbol{\sigma}\right) \quad (8)$$

while for a pure Lorentz boost B of rapidity r in the direction \mathbf{n}

$$M_{\mathbf{2}}(B) = \exp\left(-\frac{1}{2}r\mathbf{n} \cdot \boldsymbol{\sigma}\right), \quad M_{\bar{\mathbf{2}}}(B) = \exp\left(+\frac{1}{2}r\mathbf{n} \cdot \boldsymbol{\sigma}\right). \quad (9)$$

(b) The more familiar β and γ parameters of a boost are related to its rapidity r as

$$\beta = \tanh(r), \quad \gamma = \cosh(r), \quad \beta\gamma = \sinh(r). \quad (10)$$

Show that in terms of these parameters, eqs. (9) translate to

$$M_{\mathbf{2}}(B) = \sqrt{\gamma} \times \sqrt{1 - \beta\mathbf{n} \cdot \boldsymbol{\sigma}}, \quad M_{\bar{\mathbf{2}}}(B) = \sqrt{\gamma} \times \sqrt{1 + \beta\mathbf{n} \cdot \boldsymbol{\sigma}}. \quad (11)$$

(c) Let $M = M_{\mathbf{2}}(L)$ and $\bar{M} = M_{\bar{\mathbf{2}}}(L)$ be matrices representing the same continuous Lorentz symmetry $L \in SO^+(3, 1)$ in the $\mathbf{2}$ and the $\bar{\mathbf{2}}$ spinor representations. Use

eqs. (8) and (9) to show that for any such L ,

$$\overline{M} = \sigma_2 M^* \sigma_2 \quad \text{and} \quad M = \sigma_2 \overline{M}^* \sigma_2. \quad (12)$$

Hint: prove and use $\sigma_2 \boldsymbol{\sigma}^* \sigma_2 = -\boldsymbol{\sigma}$.

Next, consider the vector representation of the Lorentz symmetry and the equivalent bi-spinor representation of the $SL(2, \mathbf{C})$. In the matrix form, the $(j_+ = j_- = \frac{1}{2})$ bi-spinor multiplet of $SL(2, \mathbf{C})$ is a complex 2×2 matrix V which transforms according to

$$V \mapsto V' = M \times V \times M^\dagger \quad \text{for } M \in SL(2, \mathbf{C}). \quad (13)$$

Let's identify this bi-spinor with a Lorentz vector V^μ according to

$$V = V^\mu \overline{\sigma}_\mu = V^0 \mathbf{1}_{2 \times 2} + \mathbf{V} \cdot \boldsymbol{\sigma}, \quad (14)$$

where I follow the Peskin&Schroeder convention for the 2×2 matrices σ^μ and $\overline{\sigma}^\mu$:

$$\sigma^\mu = \overline{\sigma}_\mu = (1_{2 \times 2}, +\boldsymbol{\sigma}), \quad \overline{\sigma}^\mu = \sigma_\mu = (1_{2 \times 2}, -\boldsymbol{\sigma}). \quad (15)$$

The bi-spinor transform (13) defines a linear transform

$$V^\mu \mapsto V'^\mu = L^\mu_\nu V^\nu \quad (16)$$

of the vector V^μ .

- (d) Show that this transform is real (real V'^μ for real V^ν) and Lorentzian (preserves $V'^\mu V'_\mu = V^\nu V_\nu$). Hint: show that $\det(V) = V_\mu V^\mu$.
- (e) Show that the Lorentz transform (16) is orthochronous.
For extra challenge, show that it is also proper ($\det(L) = +1$) and therefore continuous, $L \in SO^+(3, 1)$.
- (f) Verify that this $SL(2, \mathbf{C}) \rightarrow SO^+(3, 1)$ map respects the group law, $L^\mu_\nu(M_2 M_1) = L^\mu_\lambda(M_2) L^\lambda_\nu(M_1)$.

Finally, consider the tensor representations of the Lorentz symmetry.

- (g) Show that the $(j_+ = 1, j_- = 1)$ representation is equivalent to a 2-index symmetric traceless tensor, $T^{\mu\nu} = T^{\nu\mu}$, $g_{\mu\nu}T^{\mu\nu} = 0$.

Also, show that the reducible $(j_+ = 1, j_- = 0) + (j_+ = 0, j_- = 1)$ representation is equivalent to a 2-index antisymmetric tensor, $F^{\mu\nu} = -F^{\nu\mu}$.

Hint: For any kind of angular momentum — Hermitian or not, — the tensor product of two doublets is a triplet plus a singlet, $(j = \frac{1}{2}) \otimes (j = \frac{1}{2}) = (j = 1) \oplus (j = 0)$.

3. Next, an exercise in Dirac matrices γ^μ . In this problem, you should not assume any explicit matrices for the γ^μ but simply use the anticommutation relations

$$\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = 2g^{\mu\nu}. \quad (17)$$

When necessary, you may also assume that the Dirac matrices are 4×4 , and the γ^0 matrix is hermitian while the $\gamma^1, \gamma^2, \gamma^3$ matrices are antihermitian, $(\gamma^0)^\dagger = +\gamma^0$ while $(\gamma^i)^\dagger = -\gamma^i$ for $i = 1, 2, 3$.

- (a) The original Dirac equation used $\beta = \gamma^0$ and $\alpha^i = \gamma^0\gamma^i$ (for $i = 1, 2, 3$) instead of the γ^μ . Show that eqs. (17) are equivalent to requiring all 4 matrices β and α^i to anticommute with each other and to square to 1.

- (b) Show that $\gamma^\alpha\gamma_\alpha = 4$, $\gamma^\alpha\gamma^\nu\gamma_\alpha = -2\gamma^\nu$, $\gamma^\alpha\gamma^\mu\gamma^\nu\gamma_\alpha = 4g^{\mu\nu}$,
and $\gamma^\alpha\gamma^\lambda\gamma^\mu\gamma^\nu\gamma_\alpha = -2\gamma^\nu\gamma^\mu\gamma^\lambda$.

Hint: use $\gamma^\alpha\gamma^\nu = 2g^{\nu\alpha} - \gamma^\nu\gamma^\alpha$ repeatedly.

- (c) The electron field in the EM background obeys the *covariant Dirac equation* $(i\gamma^\mu D_\mu - m)\Psi(x) = 0$ where $D_\mu\Psi = \partial_\mu\Psi - ieA_\mu\Psi$. Show that this equation implies

$$(D_\mu D^\mu + m^2 - eF_{\mu\nu}S^{\mu\nu})\Psi(x) = 0. \quad (18)$$

Besides the 4 Dirac matrices γ^μ , there is another useful matrix $\gamma^5 \stackrel{\text{def}}{=} i\gamma^0\gamma^1\gamma^2\gamma^3$.

- (d) Show that the γ^5 anticommutes with each of the γ^μ matrices — $\gamma^5\gamma^\mu = -\gamma^\mu\gamma^5$ — and commutes with all the spin matrices, $\gamma^5 S^{\mu\nu} = +S^{\mu\nu}\gamma^5$.

(e) Show that the γ^5 is hermitian and that $(\gamma^5)^2 = 1$.

(f) Show that $\gamma^5 = (i/24)\epsilon_{\kappa\lambda\mu\nu}\gamma^\kappa\gamma^\lambda\gamma^\mu\gamma^\nu$ and that $\gamma^{[\kappa}\gamma^\lambda\gamma^\mu\gamma^\nu]} = +24i\epsilon^{\kappa\lambda\mu\nu}\gamma^5$.

(g) Show that $\gamma^{[\lambda}\gamma^\mu\gamma^\nu]} = +6i\epsilon^{\kappa\lambda\mu\nu}\gamma_\kappa\gamma^5$.

(h) Show that any 4×4 matrix Γ is a unique linear combination of the following 16 matrices: 1 , γ^μ , $\frac{1}{2}\gamma^{[\mu}\gamma^{\nu]}$, $-2iS^{\mu\nu}$, $\gamma^5\gamma^\mu$, and γ^5 .

* My conventions here are: $\epsilon^{0123} = -1$, $\epsilon_{0123} = +1$, $\gamma^{[\mu}\gamma^{\nu]} = \gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu$,
 $\gamma^{[\lambda}\gamma^\mu\gamma^\nu]} = \gamma^\lambda\gamma^\mu\gamma^\nu - \gamma^\lambda\gamma^\nu\gamma^\mu + \gamma^\mu\gamma^\nu\gamma^\lambda - \gamma^\mu\gamma^\lambda\gamma^\nu + \gamma^\nu\gamma^\lambda\gamma^\mu - \gamma^\nu\gamma^\mu\gamma^\lambda$, etc.

4. Since all the spin matrices $S^{\mu\nu}$ commute with the γ^5 , for all *continuous* Lorentz symmetries L^μ_ν their Dirac-spinor representations $M_D(L) = \exp(-\frac{i}{2}\Theta_{\alpha\beta}S^{\alpha\beta})$ are block-diagonal in the eigenbasis of the γ^5 . This makes the Dirac spinor Ψ a *reducible* multiplet of the continuous Lorentz group $SO^+(3, 1)$ — it comprises two different irreducible 2-component spinor multiplets, called the left-handed Weyl spinor ψ_L and the right-handed Weyl spinor ψ_R .

This decomposition becomes clear in the Weyl convention for the Dirac matrices where in 2×2 block notations

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix} \quad (19)$$

and σ^μ and $\bar{\sigma}^\mu$ as in the Peskin & Schroeder convention (15).

(a) Show that in the Weyl convention (19), the γ^5 matrix is diagonal, specifically

$$\gamma^5 \stackrel{\text{def}}{=} i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} -1 & 0 \\ 0 & +1 \end{pmatrix}. \quad (20)$$

(b) Write down explicitly matrices for the $S^{\mu\nu}$ matrices in the Weyl convention and show that

$$S^{\mu\nu} = \begin{pmatrix} S_L^{\mu\nu} & 0 \\ 0 & S_R^{\mu\nu} \end{pmatrix} \quad (21)$$

where $S_L^{\mu\nu} = S_{\mathbf{2}}^{\mu\nu}$ and $S_R^{\mu\nu} = S_{\bar{\mathbf{2}}}^{\mu\nu}$ are respectively the $\mathbf{2}$ and $\bar{\mathbf{2}}$ representations of the Lorentz generators.

In light of eqs. (21), the Dirac spinor is a reducible $\mathbf{2} + \overline{\mathbf{2}}$ multiplet of the Spin(3, 1) Lorentz group, and for any continuous Lorentz transform L we have

$$M_D(L) = \begin{pmatrix} M_L(L) & 0 \\ 0 & M_R(L) \end{pmatrix} \quad \text{for } M_L(L) = M_{\mathbf{2}}(L) \text{ and } M_R(L) = M_{\overline{\mathbf{2}}}(L). \quad (22)$$

Consequently, in the Weyl convention the 4-components Dirac spinor field $\Psi(x)$ splits into two 2-component Weyl spinor fields — the left-handed Weyl spinor field $\psi_L(x)$ and the right-handed Weyl spinor field $\psi_R(x)$ — which transform independently (from each other) under the continuous Lorentz symmetries,

$$\Psi_{\text{Dirac}}(x) = \begin{pmatrix} \psi_L(x) \\ \psi_R(x) \end{pmatrix} \quad \text{where} \quad \begin{aligned} \psi'_L(x') &= M_L(L)\psi_L(x), \\ \psi'_R(x') &= M_R(L)\psi_R(x). \end{aligned} \quad (23)$$

- (c) Use eqs. (12) to show that the hermitian conjugate of each Weyl spinor transforms equivalently to the other spinor. Specifically, the $\sigma_2 \times \psi_L^*(x)$ transforms under continuous Lorentz symmetries like the $\psi_R(x)$, while the $\sigma_2 \times \psi_R^*(x)$ transforms like the $\psi_L(x)$.

Note: the * superscript on a multi-component quantum field means hermitian conjugation of each component field but without transposing the components, thus

$$\psi_L = \begin{pmatrix} \psi_{L1} \\ \psi_{L2} \end{pmatrix}, \quad \psi_L^* = \begin{pmatrix} \psi_{L1}^\dagger \\ \psi_{L2}^\dagger \end{pmatrix}, \quad \text{while} \quad \psi_L^\dagger = (\psi_{L1}^\dagger \quad \psi_{L2}^\dagger), \quad (24)$$

and likewise for the ψ_R and its conjugates.

Next, consider the Dirac Lagrangian $\mathcal{L} = \overline{\Psi}(i\gamma^\mu\partial_\mu - m)\Psi$.

- (d) Express this Lagrangian in terms of the Weyl spinor fields $\psi_L(x)$ and $\psi_R(x)$ (and their conjugates $\psi_L^\dagger(x)$ and $\psi_R^\dagger(x)$).
- (e) Show that for $m = 0$ — and only for $m = 0$ — the two Weyl spinor fields become independent from each other.