

This homework has 4 problems. Problems 1 and 2 are about non-abelian gauge theories. Problems 3 and 4 are about the Lorentz symmetry and its generators. Altogether, it's a pretty large homework set, so start working early.

1. In class, I have focused on the *fundamental multiplet* of the local $SU(N)$ symmetry, *i.e.*, a set of N fields (complex scalars or Dirac fermions) which transform as a complex N -vector,

$$\Psi'(x) = U(x)\Psi(x) \quad i.e. \quad \Psi'^i(x) = \sum_j U^i_j(x)\Psi^j(x), \quad i, j = 1, 2, \dots, N \quad (1)$$

where $U(x)$ is an x -dependent unitary $N \times N$ matrix, $\det U(x) \equiv 1$. Now consider $N^2 - 1$ real fields $\Phi^a(x)$ forming an *adjoint multiplet*: In matrix form

$$\Phi(x) = \sum_a \Phi^a(x) \times \frac{\lambda^a}{2} \quad (2)$$

is a traceless hermitian $N \times N$ matrix which transforms under the local $SU(N)$ symmetry as

$$\Phi'(x) = U(x)\Phi(x)U^\dagger(x). \quad (3)$$

Note that this transformation law preserves the $\Phi^\dagger = \Phi$ and the $\text{tr}(\Phi) = 0$ conditions.

The covariant derivatives D_μ act on an adjoint multiplet according to

$$D_\mu\Phi(x) = \partial_\mu\Phi(x) + i[\mathcal{A}_\mu(x), \Phi(x)] \equiv \partial_\mu\Phi(x) + i\mathcal{A}_\mu(x)\Phi(x) - i\Phi(x)\mathcal{A}_\mu(x), \quad (4)$$

or in components

$$D_\mu\Phi^a(x) = \partial_\mu\Phi_a(x) - f^{abc}\mathcal{A}_\mu^b(x)\Phi^c(x). \quad (5)$$

- (a) Verify that these derivatives are indeed covariant under the finite gauge transforms (3).

- (b) Verify the Leibniz rule for the covariant derivatives of matrix products. Let $\Phi(x)$ and $\Xi(x)$ be two adjoint multiplets while $\Psi(x)$ is a fundamental multiplet and $\Psi^\dagger(x)$ is its hermitian conjugate (row vector of Ψ_i^*). Show that

$$\begin{aligned} D_\mu(\Phi\Xi) &= (D_\mu\Phi)\Xi + \Phi(D_\mu\Xi), \\ D_\mu(\Phi\Psi) &= (D_\mu\Phi)\Psi + \Phi(D_\mu\Psi), \\ D_\mu(\Psi^\dagger\Xi) &= (D_\mu\Psi^\dagger)\Xi + \Psi^\dagger(D_\mu\Xi). \end{aligned} \tag{6}$$

- (c) Show that for an adjoint multiplet $\Phi(x)$,

$$[D_\mu, D_\nu]\Phi(x) = i[\mathcal{F}_{\mu\nu}(x), \Phi(x)] = ig[F_{\mu\nu}(x), \Phi(x)] \tag{7}$$

or in components $[D_\mu, D_\nu]\Phi^a(x) = -gf^{abc}F_{\mu\nu}^b(x)\Phi^c(x)$.

- In my notations A_μ and $F_{\mu\nu}$ are the canonically normalized potential and tension fields, while $\mathcal{A}_\mu = gA_\mu$ is the connection in the covariant derivative and $\mathcal{F}_{\mu\nu} = gF_{\mu\nu}$ is the curvature of that connection.

In class, I have argued (using covariant derivatives) that the tension fields $\mathcal{F}_{\mu\nu}(x)$ themselves transform according to eq. (3). In other words, the $\mathcal{F}_{\mu\nu}^a(x)$ form an adjoint multiplet of the $SU(N)$ symmetry group.

- (d) Verify the $\mathcal{F}'_{\mu\nu}(x) = U(x)\mathcal{F}_{\mu\nu}(x)U^\dagger(x)$ transformation law directly from the definition $\mathcal{F}_{\mu\nu} \stackrel{\text{def}}{=} \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu]$ and the non-abelian gauge transform of the \mathcal{A}_μ fields.
- (e) Verify the covariant differential identity for the non-abelian tension fields $\mathcal{F}_{\mu\nu}(x)$:

$$D_\lambda\mathcal{F}_{\mu\nu} + D_\mu\mathcal{F}_{\nu\lambda} + D_\nu\mathcal{F}_{\lambda\mu} = 0. \tag{8}$$

Note the covariant derivatives (4) in this equation.

Finally, consider the $SU(N)$ Yang–Mills theory — the non-abelian gauge theory that does not have any fields except $\mathcal{A}^a(x)$ and $\mathcal{F}^a(x)$; its Lagrangian is

$$\mathcal{L}_{\text{YM}} = -\frac{1}{2g^2} \text{tr}(\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu}) = -\frac{1}{4} \sum_a F_{\mu\nu}^a F^{a\mu\nu}. \tag{9}$$

- (f) Show that the Euler–Lagrange field equations for the Yang–Mills theory can be written in covariant form as $D_\mu\mathcal{F}^{\mu\nu} = 0$.

Hint: first show that for an infinitesimal variation $\delta\mathcal{A}_\mu(x)$ of the non-abelian gauge fields, the tension fields vary according to $\delta\mathcal{F}_{\mu\nu}(x) = D_\mu\delta\mathcal{A}_\nu(x) - D_\nu\delta\mathcal{A}_\mu(x)$.

2. Continuing the previous problem, consider an $SU(N)$ gauge theory in which $N^2 - 1$ vector fields $A_\mu^a(x)$ interact with some “matter” fields $\phi_\alpha(x)$,

$$\mathcal{L} = -\frac{1}{2g^2} \text{tr}(\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu}) + \mathcal{L}_{\text{mat}}(\phi, D_\mu\phi). \quad (10)$$

For the moment, let me keep the matter fields completely generic — they can be scalars, or vectors, or spinors, or whatever, and form any kind of a multiplet of the local $SU(N)$ symmetry as long as such multiplet is complete and non-trivial. All we need to know right now is that there are well-defined covariant derivatives $D_\mu\phi$ that depend on the gauge fields A_μ^a , which give rise to the currents

$$J^{a\mu} = -\frac{\partial\mathcal{L}_{\text{mat}}}{\partial A_\mu^a} = -\sum_\phi \frac{\partial\mathcal{L}_{\text{mat}}}{\partial(D_\mu\phi)} \times ig\hat{T}^a\phi. \quad (11)$$

Collectively, these $N^2 - 1$ currents should form an adjoint multiplet of the local $SU(N)$ symmetry, meaning

$$J_\mu(x) \stackrel{\text{def}}{=} \sum_a J_\mu^a(x) \frac{\lambda^a}{2} \text{ transforms to } J'_\mu(x) = U(x)J_\mu(x)U^\dagger(x). \quad (12)$$

- (a) Show that in this theory the equation of motion for the A_μ^a fields are $D_\mu F^{a\mu\nu} = J^{a\nu}$ and that consistency of these equations requires the currents to be *covariantly conserved*,

$$D_\mu J^\mu = \partial_\mu J^\mu + i[\mathcal{A}_\mu, J^\mu] = 0, \quad (13)$$

or in components, $\partial_\mu J^{a\mu} - f^{abc}A_\mu^b J^{c\mu} = 0$.

Note: a covariantly conserved current does *not* lead to a conserved charge,

$$\frac{d}{dt} \int d^3\mathbf{x} J^{a0}(\mathbf{x}, t) \neq 0. \quad (14)$$

Now consider a simple example of matter fields — a fundamental multiplet $\Psi(x)$ of N scalar fields $\Psi^i(x)$, with a Lagrangian

$$\mathcal{L}_{\text{mat}} = D_\mu \Psi^\dagger D^\mu \Psi - m^2 \Psi^\dagger \Psi - \frac{\lambda}{4} (\Psi^\dagger \Psi)^2, \quad \mathcal{L}_{\text{net}} = \mathcal{L}_{\text{mat}} - \frac{1}{2g^2} \text{tr}(\mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu}), \quad (15)$$

and hence the $SU(N)$ currents are

$$J^{a\mu} = -g \text{Im} \left(\Psi^\dagger \lambda^a D^\mu \Psi \right) = -\frac{ig}{2} (D^\mu \Psi^\dagger) \lambda^a \Psi + \frac{ig}{2} \Psi^\dagger \lambda^a D^\mu \Psi. \quad (16)$$

(b) Derive these $SU(N)$ currents. Then use the matrix identity

$$\sum_a (\lambda^a)^i_j (\lambda^a)^k_\ell = 2\delta^i_\ell \delta^k_j - \frac{2}{N} \delta^i_j \delta^k_\ell \quad (17)$$

to show that in the matrix form

$$\begin{aligned} J_\mu &\stackrel{\text{def}}{=} \sum_a J_\mu^a \times \frac{1}{2} \lambda^a = \frac{ig}{2} \left((D_\mu \Psi) \otimes \Psi^\dagger - \Psi \otimes D_\mu \Psi^\dagger \right) \\ &\quad - \frac{ig}{2N} \left(\Psi^\dagger D_\mu \Psi - (D_\mu \Psi^\dagger) \Psi \right) \times \mathbf{1}_{N \times N} \end{aligned} \quad (18)$$

where $(D_\mu \Psi) \otimes \Psi^\dagger$ denotes $N \times N$ matrix with elements $(D_\mu \Psi)^i \times \Psi_j^*$, and likewise for the $\Psi \otimes D_\mu \Psi^\dagger$.

(c) Verify that under the $SU(N)$ gauge transforms, the currents (16) transform into each other as members of the adjoint multiplet, *i.e.*, the matrix (18) transforms according to eq. (12).

(d) Finally, verify the covariant conservation $D_\mu J^{a\mu}$ of these currents when the scalar fields $\Psi^i(x)$ and $\Psi_i^\dagger(x)$ obey their equations of motion.

3. Now consider a different subject, namely the continuous Lorentz group $SO^+(3,1)$ and its generators $\hat{J}^{\mu\nu} = -\hat{J}^{\nu\mu}$. In 3D terms, the six independent $\hat{J}^{\mu\nu}$ generators comprise the 3 components of the angular momentum $\hat{J}^i = \frac{1}{2}\epsilon^{ijk}\hat{J}^{jk}$ — which generate the rotations of space — plus 3 generators $\hat{K}^i = \hat{J}^{0i} = -\hat{J}^{i0}$ of the Lorentz boosts.

(a) In 4D terms, the commutation relations of the Lorentz generators are

$$\left[\hat{J}^{\alpha\beta}, \hat{J}^{\mu\nu}\right] = ig^{\beta\mu}\hat{J}^{\alpha\nu} - ig^{\alpha\mu}\hat{J}^{\beta\nu} - ig^{\beta\nu}\hat{J}^{\alpha\mu} + ig^{\alpha\nu}\hat{J}^{\beta\mu}. \quad (19)$$

Show that in 3D terms, these relations become

$$\left[\hat{J}^i, \hat{J}^j\right] = i\epsilon^{ijk}\hat{J}^k, \quad \left[\hat{J}^i, \hat{K}^j\right] = i\epsilon^{ijk}\hat{K}^k, \quad \left[\hat{K}^i, \hat{K}^j\right] = -i\epsilon^{ijk}\hat{J}^k. \quad (20)$$

The Lorentz symmetry dictates the commutation relations of the $\hat{J}^{\mu\nu}$ with any operators comprising a Lorentz multiplet. In particular, for any Lorentz vector \hat{V}^μ

$$\left[\hat{V}^\lambda, \hat{J}^{\mu\nu}\right] = ig^{\lambda\mu}\hat{V}^\nu - ig^{\lambda\nu}\hat{V}^\mu. \quad (21)$$

- (b) Spell out these commutation relations in 3D terms, then use them to show that the Lorentz boost generators $\hat{\mathbf{K}}$ do not commute with the Hamiltonian \hat{H} .
- (c) Show that even in the non-relativistic limit, the Galilean boosts $t' = t$, $\mathbf{x}' = \mathbf{x} + \mathbf{v}t$ and their generators $\hat{\mathbf{K}}_G$ do not commute with the Hamiltonian operator of a QFT or a quantum system of several particles.

Note: Only the *time-independent* symmetries commute with the Hamiltonian. But when the action of a symmetry is manifestly time dependent — like a Galilean boost $\mathbf{x}' = \mathbf{x} + \mathbf{v}t$, or a Lorentz boost — the symmetry operators do not commute with the time evolution and hence with the Hamiltonian.

4. Next, consider the little group $G(p)$ of Lorentz symmetries preserving some momentum 4-vector p^μ . For the moment, allow the p^μ to be time-like, light-like, or even space-like — anything goes as long as $p \neq 0$.

(a) Show that the little group $G(p)$ is generated by the 3 components of the vector

$$\hat{\mathbf{R}} = p^0 \hat{\mathbf{J}} + \mathbf{p} \times \hat{\mathbf{K}} \quad (22)$$

after a suitable component-by-component rescaling.

Suppose the momentum p^μ belongs to a massive particle, thus $p^\mu p_\mu = m^2 > 0$. For simplicity, assume the particle moves in z direction with velocity β , thus $p^\mu = (E, 0, 0, p)$ for $E = \gamma m$ and $p = \beta \gamma m$. In this case, the properly normalized generators of the little group $G(p)$ are the

$$\begin{aligned} \tilde{J}^x &= \frac{1}{m} \hat{R}^x = \gamma \hat{J}^x - \beta \gamma \hat{K}^y, \\ \tilde{J}^y &= \frac{1}{m} \hat{R}^y = \gamma \hat{J}^y + \beta \gamma \hat{K}^x, \\ \tilde{J}^z &= \frac{1}{\gamma m} \hat{R}^z = \hat{J}^z, \quad \text{the helicity.} \end{aligned} \quad (23)$$

(b) Show that these generators have angular-momentum-like commutators with each other, $[\tilde{J}^i, \tilde{J}^j] = i\epsilon^{ijk} \tilde{J}^k$. Consequently, the little group $G(p)$ is isomorphic to the rotation group $SO(3)$.

Now suppose the momentum p^μ belongs to a massless particle, $p^\mu p_\mu = 0$. Again, assume for simplicity that the particle moves in the z direction, thus $p^\mu = (E, 0, 0, E)$. In this case, we cannot normalize the generators of the little group as in eq. (23); instead, let's normalize them according to

$$\hat{\mathbf{I}} = \frac{1}{E} \hat{\mathbf{R}} = \hat{\mathbf{J}} + \vec{\beta} \times \hat{\mathbf{K}}, \quad (24)$$

or in components,

$$\hat{I}^x = \hat{J}^x - \hat{K}^y, \quad \hat{I}^y = \hat{J}^y + \hat{K}^x, \quad \hat{I}^z = \hat{J}^z. \quad (25)$$

(c) Show that these generators obey similar commutation relations to the \hat{p}^x , \hat{p}^y , and \hat{J}^z

operators, namely

$$[\hat{J}^z, \hat{I}^x] = +i\hat{I}^y, \quad [\hat{J}^z, \hat{I}^y] = -i\hat{I}^x, \quad [\hat{I}^x, \hat{I}^y] = 0. \quad (26)$$

Consequently, the little group $G(p)$ is isomorphic to the ISO(2) group of *rotations and translations* in the xy plane.

- (d) Finally, show that for a tachyonic momentum with $p^\mu p_\mu < 0$, the properly normalized generators of the little group have similar commutation relations to the \hat{K}^x , \hat{K}^y , and \hat{J}^z operators. Consequently, the little group $G(p)$ is isomorphic to the $SO^+(2,1)$, the continuous Lorentz group in $2 + 1$ spacetime dimensions.